

Distribution and abundance of benthic macroorganisms in and around Visakhapatnam Harbour on the east coast of India

Running head: *Distribution and abundance of benthic macroorganism*

Amar S Musale, Dattesh V Desai*, S S Sawant, K Venkat, A C Anil

CSIR-National Institute of Oceanography, Dona Paula, Goa.

Phone (office) 91(0)832-2450512; Fax: 91(0)832-2450615.

*Corresponding author, E-mail: ddattesh@nio.org

ABSTRACT

Benthic communities form an important component of the marine food chain. Their occurrence also provides information on the health of the ecosystem. A study was carried out to understand the distribution and abundance of macrobenthos along with sediment characteristics and physicochemical parameters in Visakhapatnam harbour, a major port along the east coast of India. In all 84 macrobenthic taxa were reported from the port area of which 60 were polychaetes and 24 were other invertebrate taxa. Our observations revealed an increase in the number of polychaete species observed over the last 20 years from this region. An earlier study reported 38 polychaete species in 1975 and a year later the number of polychaete species reported were 12, indicating an increase in the number of polychaete species in the present study to about 150%. The macrobenthic abundance and dominance of species varied with the seasons. Pre-monsoon was dominated by Cirratulus sp., during monsoon tanaids were dominant indicating a seasonal shift in the occurrence and dominance of macrobenthos. During post-monsoon, Cossura coasta was dominant followed by Nephtys dibranchis and amphipods. Sediment characteristics (sand, silt and clay), organic carbon and dissolved oxygen were the important factors to influence the abundance and species diversity. The abundance of macrobenthic forms also varied with inner and outer harbour region. Higher species diversity was observed in the outer harbour suggesting outer harbour to have semi-polluted conditions such as higher dissolved oxygen (DO) and salinity, low nutrients (nitrite, nitrate and silicate), and low organic carbon of the sediment.

Keywords: Macrobenthos, Polychaete, Visakhapatnam Harbour, Sediment, Organic carbon

INTRODUCTION

Estuaries and coasts are continuously disturbed by both natural and anthropogenic activities. Harbours located in estuaries are in particular often subjected to various forms anthropogenic pressure. These include sewage or municipal runoff, terrestrial runoff during the monsoon, and port related activities such as dredging, oil spill, petroleum effluents, out-fall of variety of cargo handled by the port etc. all of which disturb the port environment. Visakhapatnam is a major harbour in the North East of Andhra Pradesh, on the east coast of India (Latitude: 17° 41' N and Longitude: 83 ° 18' E). The macrobenthic community of the Visakhapatnam harbour was previously studied about two decades ago to understand the impact of pollution in the harbour area (Ganapati, 1969; Ganapati & Raman, 1979; Raman & Ganapati, 1983; Raman, 1995). The present investigation was carried under a port biological baseline survey as a part of ballast water management program.

Visakhapatnam harbour is known to be highly polluted due to discharge of industrial effluents and domestic sewage (Ganapati & Raman, 1973; Raman, 1995). The land-locked situation (where there is a narrow entrance channel which forms the main outlet for the harbour waters into the Bay of Bengal) of the harbour and the limitations of natural flushing processes have largely contributed to exposure of flora and fauna to environmental pollution stress from effluents of a variety of industrial installations which have sprung up in the environs of harbour in the recent years (Ganapati & Raman 1973,1979; Sarma et al., 1982; Raman & Ganapati, 1986; Rathod et al., 1995; Kalavati etl al., 1997; Tripathy et al., 2005). The harbour also receives most of the city's untreated domestic effluent which is considerable owing to rapidly growing urban population (Raman, 1995). The earlier studies have pointed out higher pollution in the inner harbour compared to the outer harbour. A observation carried out between 1985-87 indicated that the nitrite levels were 0-0.7 mg/l; nitrate at 0.6-6.16 mg/l; phosphate at 0-8.07mg/l; and silicate levels were 0.16 – 3.2 mg/l. Thus it is important to evaluate the present condition of port environment by comparing the nutrients levels which were reported earlier and their influence on the population of macrobenthos. There is also considerable freshwater runoff into the harbour through a monsoon-fed stream known as 'Mehadrigedda' with annual mean discharge of approx. $2.1 \text{ m}^3 \text{ s}^{-1}$ (Raman & Ganapati, 1983).

Macrobenthic polychaetes inhabit the sediment surface and are dependent upon the benthic sediment characteristics and also on the near bottom water quality. Some polychaetes (e.g. *Capitella capitata*, *Nereis glandisincta*, *Diopatra neapolitana*, *Nephtys oligobranchia*) live in conditions of extremely high levels of organic content and also low levels of dissolved oxygen. They have been considered as the indicators of organic pollution. Since there is a gap of nearly two decades on the data available

on the macrobenthos population with respect to ever growing industrial growth, increase in the port related activities and also after the construction of outer harbor which was completed in 1975, only one report on ecology of macrobenthos from this region is available 1986 (Bismillah, 1986). Increasing pollution and sluggish circulation will lead to increase in nutrient accumulation and regeneration and this might favor some of the pollution tolerant polychaete species and needs to be studied in detail. In this study we investigated the distribution of macrobenthos in the harbour along with the quality of the environment in terms of physico-chemical parameters and nutrients, which will also aid in commenting on the health of the harbour. Since, inner harbour experiences more pollution and less flushing compared to the outer harbour, it is important to conduct the study to identify spatial and temporal variation in the macrobenthic population. Efforts were also made to find out the relation between sediment characteristics and macrobenthic community in and around the harbour. The study during different seasons will provide information how different seasons affect the environment and in turn the macrobenthos population in the Visakhapatnam harbour, especially during non-monsoon months when the salinity will be higher where in, more number of euryhaline forms can proliferate. However, with the onset of monsoon and with an increase in the fresh water input into the harbour might bring drastic change in salinity (as low as 4.6 psu) (Raman & Ganapati, 1983) and other abiotic factors (dissolved oxygen and nutrients). The land runoff fills the harbour area with organic material which can enhance the organic carbon levels. This can lead to an accumulation of toxic material which can change or reduce the sediment quality and macrobenthic abundance. There is inadequate tidal flushing and stagnant conditions prevailing in Visakhapatnam harbour (Sarma *et al.*, 1982; Raman, 1995) and only during monsoon the IH gets flushed out due to land runoff. In the present study we would like to put forward the following objectives (a) To evaluate the spatio-temporal variation in the macrobenthos population (b) Spatio-temporal variation in the levels of nutrients and other abiotic parameters (c) Implication of environmental parameters (water parameters and sediment characteristics) on the macrobenthos population (d) to provide a comparative account of macrobenthos population with respect to earlier reports in the Visakhapatnam harbour.

MATERIALS AND METHODS

Description of the study area

Visakhapatnam harbour is a semi-enclosed water body situated in north-east of Andhra Pradesh state, east coast of India (17° 41'N and 83° 18' E). The harbour area extends 4km from north-western arm of inner to the outer area of the harbour (RamaRaju *et al.*, 1990). The inner harbour (subsequently

abbreviated as IH) area is land-locked area which consists of naturally protected entrance channel, turning basin and three navigable arms (northern, western and north-western arm). In contrast, the outer harbour (subsequently abbreviated as OH) area has access to the open sea (Bay of Bengal) through the entrance channel and has a protected basin with two breakwaters. There are 18 and 6 berths in the IH and OH area with two moorings respectively, indicating the amount of cargo handled in the inner harbour will be more. The climate in Visakhapatnam is governed by its location in the tropics which is mainly affected by seasonal monsoon (south-west during June to September and north-east, October to December). Visakhapatnam receives around 1000mm of rainfall on an average annually, out of which south-west accounts to 70% and remaining by north-east monsoon with few showers during rest of the year. Wave height is also highest during south-west monsoon compared to other seasons (Suresh et al., 2012). Such a difference in the pattern of monsoon will have impact on the sea water characteristics and terrestrial runoff into the harbour.

A total of 24 sampling stations were selected in and around the Visakhapatnam harbour (Fig. 1). 13 stations were selected in Inner Harbour namely DC-Jetty (IH-1), Boat basin berth-3 (IH-2), Port dry dock (IH-3), Turning basin (IH-4), Hindustan shipyard (IH-5), Oil refinery berth (IH-6), Fertilizer wharf (IH-7), East quay berth-1 (IH-8), West quay berth-1 (IH-9), West quay berth-4 (IH-10), East quay berth-5 (IH-11), East quay berth-7 (IH-12), East quay berth-9 (IH-13) and 11 stations were selected in Outer Harbour namely Dredger berth (OH-1), General cargo berth-S (OH-2), General cargo berth-N (OH-3), Ore berth-1 (OH-4), Ore berth-2 (OH-5), Container berth-1 (OH-6), Container berth-2 (OH-7), Fishery jetty (OH-8), Oil berth (OH-9), Turning circle (OH-10) and LPG berth (OH-11) (Fig.1). Sampling was carried out during January 2007 (First post monsoon) and December 2007 (Second post monsoon; north-east monsoon, November-December 2007), April 2008 (Pre-monsoon) and August 2008 (south-west monsoon), representing four different seasons. Near bottom water samples were collected using Niskin water sampler. Water samples were analyzed for salinity, dissolved oxygen (DO), temperature and nutrients (nitrite, nitrate, phosphate and silicate) following the methods described by Parsons *et al.* (1984). The samples for sea water analysis were collected in triplicate. Sediment samples were collected from an average depth of 8-10 m using van Veen grab (0.04m²). At each station three grab samples were taken (n=3). These samples were washed separately through a 500µm nylon mesh at sea and then transferred to polythene bags and preserved in 5% formaldehyde in sea water containing rose bengal stain. In the laboratory sediment samples were sieved through a 500µm metal sieve and all macrobenthic fauna were preserved in plastic vials containing 5% formaldehyde solution for further microscopic analysis. Polychaetes were identified to the highest taxonomic resolution (genus or species) possible

with the help of available identification keys (Day, 1967; Gosner, 1971). The polychaetes recorded were provided with following species codes (*Nephtys dibranchis*, NepD; *Nephtys capensis*, NepC; *Flabelligera* sp., FlabS; *Nephtys polybranchia*, NepP; *Orbinella* sp., OrbS; *Nephtys* sp., NepS; *Nereis* sp., NerS; *Diopatra neopolitana*, DioN; *Hesionella* sp., HesS; *Diopatra* sp., DioS; *Syllis* sp., SylS; *Cossura coasta*, CosC; *Sabella* sp., SabS; *Cossura* sp., CosS; *Aricidea* sp., AriS; *Prionospio pinnata*, PriP; *Ampharete* sp., AmpS; *Prionospio cirrifera*, PriCfr; *Chone* sp., ChoS; *Prionospio cirrobranchiata*, PriCbr; *Leocrates* sp.(Hesionidae), LeoS; *Prionospio* sp., PriS; *Phyllodoce* sp., PhyS; *Ancistrosyllis constricta*, AnsC; *Paraonis* sp., ParS; *Ancistrosyllis* sp., AnsS; *Sternaspis scutata*, SterS; *Cirratulus chrysoderama*, CirChr; *Amphiglena mediterranea*, AmplM; *Cirratulus concinnus*, CirCon; *Boccardia* sp., BocS; *Cirratulus cirratus*, CirCir; *Dorvillea* sp., DorS; *Cirratulus filiformis*, CirF; *Serpula* sp., SerS; *Cirratulus* sp., CirS; *Maldane* sp., MalS; *Capitella capitata*, CapC; *Levensenia* sp., LevS; *Capitella* sp., CapS; *Heteromastus filiformis*, HetF; *Heteromastus* sp., HetS; *Mediomastus capensis*, MedC; *Mediomastus* sp., MedS; *Magelona rosea*, MagR; *Magelona cincta*, MagC; *Magelona* sp., MagS; *Glycera alba*, GlyA; *Glycera* sp., GlyS; *Poecilochaetus serpen*, PoeSr; *Poecilochaetus* sp., PoeS; *Glycinde capensis*, GlyC; *Scoloplos chevalieri*, ScoC; *Scoloplos* sp., ScoS; *Polydora* sp., PolS; *Onuphis eremita*, OnuE; *Onuphis* sp., OnuS; *Pisione oerstedii*, PisO; *Euclymene* sp., EuclS; *Pseudoeurythoe* sp., PseuS). Other macro-fauna (non-polychaete) belonging to crustaceans, molluscs, echinoderms, oligochaetes, nemerteans, sipunculans and fish larvae were identified to the group level. Numerical abundance of each species was recorded and expressed as number of species per square meter (no. m⁻²). Organic carbon (OC) and percentage composition of sediment (sand, silt and clay) were determined by standard titration methods and pipette analysis respectively (Wakeel & Riley, 1956; Buchanan, 1984). Organic carbon was expressed as percentage of sediment dry weight.

Macrobenthic fauna, especially polychaetes, reflect the ecological and environmental status and this was explained in terms of number of individuals or specimens (N), number of species (S), total abundance (A), Margalef species richness (d), Pielou's evenness (J') and Shannon index (H') using log₂ scale at each station (Clarke & Gorley, 2001). Bray-Curtis similarity for species diversity for macrobenthic polychaetes was determined using PRIMER-v5 (Clarke & Gorley, 2011). Seasonal variation in the total macrobenthic community, polychaetes and other invertebrate taxa is presented using SURFER-6 (developed by Golden Software Inc., USA). Canonical correspondence analysis (CCA) was performed to evaluate the relationship between environmental variables and macrobenthic polychaetes as well as other taxonomic groups (ter Braak, 1995) using the Multi-Variate Statistical Package version 3.1 (Kovach, 1998).

RESULTS

Sediment characteristics

It was observed that in all four seasons (first post monsoon, second post monsoon, pre-monsoon and monsoon), sediments were mainly of four types viz: sand, silt, clay and mixed (constituting almost equal proportions of sand, silt and clay) (Fig. 2a-d). IH was dominated by sand and OH was dominated by silt during first post monsoon season (Fig. 2a). Whereas, during second post monsoon season silt was higher at IH (Fig. 2b). However, pre-monsoon season was distinguished by silt and sand dominating at both inner and outer harbours (Fig. 2c). Mixed type of sediment was observed during monsoon season at IH wherein sand was dominated at four stations and in OH sand and silt dominated the sediment (Fig. 2d).

Sediment organic carbon ranged from 0.36 to 3.56% in IH, while at OH it ranged between 0.29-3.47% during first post monsoon (Fig. 2a). During second post monsoon organic carbon ranged from 0.5 to 3.8% in IH and at OH it ranged from 0.4 to 3.8% (Fig. 2b). During pre-monsoon season, organic carbon was high at IH when compared to OH which indicated a large variation (0.53% at Ore berth-2; OH5 to 3.6% at Ore berth-1; OH-4) (Fig. 2c). Monsoon season showed higher organic carbon (0.92-4.6% in IH and 0.7-4.6% in OH) when compared to other seasons (Fig. 2d).

Hydrological parameters

Temperature was high during the monsoon and ranged between 28 to 29.8°C. Temperature it was lowest during post monsoon (25.3 to 228.0°C). During the pre-monsoon season temperature ranged from 27.6 to 28.7°C (Table 2). Salinity was high during pre-monsoon (ranging from 30.1 to 31.9psu) but minimum salinity occurred during the monsoon (ranging from 20.2 to 28.2psu). First post monsoon showed higher salinity (29.0 to 31.1psu) compared to second post monsoon season (23.8 to 26psu) (Table 2). Dissolved oxygen (DO) showed very high inter-seasonal variations. During first post monsoon season DO ranged from 2.3 to 6.2 mg. L⁻¹ and during second post monsoon season it was 2.9 to 6.0 mg. L⁻¹. However, during pre-monsoon and monsoon the DO ranged from 1 to 6.1 mg. L⁻¹ and 0.7 to 5.9 mg. L⁻¹ respectively, indicating the lower levels to be hypoxic (Table 2).

Nutrients

During pre-monsoon season nitrite concentration was low (1.8 µM) in OH compared to IH (Table 2). There was a large difference in the nitrate values in IH (6.1 to 132.0 µM) compared to OH (0.7 to 39.8 µM) during post monsoon II (northeast monsoon). Similar was the case during pre-monsoon also where nitrate concentration was high in IH. During monsoon maximum nitrate concentration at

IH was 21.0 μM and at OH it was 6.1 μM . Concentration of phosphate and silicate was high in IH compared to OH throughout the study (Table 2).

Macrobenthic community

Figure 3 (a-d) depicts the abundance of macrobenthos at IH and OH during different seasons. The macrobenthic fauna comprised of polychaetes, crustaceans, molluscs, oligochaetes, sipunculans (peanut worms), and nemerteans (ribbon worms). Among these groups polychaetes were the most common during all the seasons. Of a total of 84 macrobenthic forms 60 were polychaetes. Polychaetes were most abundant during all the seasons except monsoon, and their contribution was more than 20% to the total macrobenthic abundance. During monsoon season tanaiids were the most dominant group that contributed 29% to the total macrobenthic abundance (Table 3).

Seasonal variation in the abundance of polychaetes

The cossurid polychaete, *Cossura coasta* was dominant during first and second post monsoon contributing 27 and 28% of total macrobenthic abundance (Table 3; Fig. 4a, b&e). However, its dominance was restricted only to the OH as in IH, *Nephtys dibranchis* (11%) was the dominant species during first post monsoon and amphipods were dominant (15%) during second post monsoon season (Table 3; Fig. 4 a&b). During pre-monsoon IH was dominated by *Cirratulus* sp., which contributed 13% to the total macrobenthic abundance. However OH during this season was dominated by *Cossura coasta* (19%) (Fig. 4c). During monsoons tanaiids dominated the macrobenthic abundance contributing 29% to the total, and they were confined only to IH (Fig. 4d). It was noted that *Cossura coasta*, *Cirratulus* sp., *Nephtys dibranchis* are the polychaete species which contributed considerably to the total macrobenthic abundance (Fig. 4e). In general, polychaetes were dominant during all the seasons except monsoon (Table 3). A significant (one way ANOVA $p \leq 0.02$) difference in the abundance of macrobenthic polychates was observed with respect to seasons.

Seasonal variation in the abundance of macrobenthos belonging to other groups

Macrobenthos belonging to crustaceans, molluscs, oligochaetes, sipunculans (peanut worms) and nemerteans (ribbon worms) were recorded during all the seasons, but their contribution to the total macrobenthic abundance varied with the seasons (Table 3; Fig. 5a-d). Amphipods (post-monsoon II–15%, pre-monsoon–8%), tanaiids (monsoon–29%), and sipunculans (post monsoon I–5%) are major contributors among the other groups to the total macrobenthos (Table 3). In general, crustaceans were abundant and commonly encountered in IH during all the seasons except first post monsoon,

when sipunculans showed dominance over the crustaceans (Table 3). However, in OH crustaceans were dominant only during second post monsoon and pre-monsoon seasons and other two seasons were dominated by nemerteans followed by bivalves (first post monsoon). Oligochaetes were dominant during monsoon season. Crustaceans were dominant among other invertebrate macrobenthic taxa mainly comprised of tanaids, amphipods, ostracods, isopods, decapods, lucifer larvae and copepods (Fig. 5a-d).

Species diversity index for polychaetes

Species diversity index at all the stations were estimated based on Margalef species richness (d), Shanon-Weiner index (H') and evenness (J'). The maximum numbers of species were encountered during first post-monsoon and monsoon season ($S=17$ at both the seasons) (Table 1a and d). The correspondence values of the Shannon-Weiner index (H') were also high during these seasons (2.45 and 2.33 during first post-monsoon and monsoon season respectively). During second post monsoon season, species diversity was low compared to other three seasons (Table 1b). In pre-monsoon season 16 species were observed (Table 1c).

Bray-Curtis similarity index was applied for grouping the stations according to polychaete abundance. At 50% similarity level, five groups were revealed during first post monsoon season (Fig.6a). The stations in the first group had high DO levels and the second group stations had high DO and low organic carbon (<3%). In the third and fourth group stations, the polychaete, *Nephtys dibranchis* was the dominant and most commonly represented species and these stations had low DO. Fifth group stations had low organic carbon (<3%). Second post monsoon season revealed three groups and six dissimilar stations. Nine stations were grouped together which had low organic carbon and *Cossura coasta* as the dominant species, whereas second group represented two stations with low DO and organic carbon. Stations with higher abundance of *Cirratulus* sp. were in third group which were associated with silty substratum and low DO (Fig. 6b). During pre-monsoon, stations with low organic carbon and high DO formed the first group. Stations with silty substratum (>62.2%) and low DO were in second (IH9 and IH4) and stations with moderate silt were in third group (OH2 and OH8) respectively. Fourth group stations were sandy substratum and high DO and fifth group was with low organic carbon (<3%) (Fig.6c). During monsoon, stations were grouped according to sediment characteristics and environmental parameters, organic carbon being a responsible factor for grouping of stations. Occurrence and abundance of *Nephtys dibranchis*, *Cossura* sp., and *Cirratulus* sp. also affected the grouping of stations (Fig.6d).

Canonical correspondence analysis (CCA) for macrobenthic polychaetes and environmental variables

In the CCA biplot for polychaetes four axis represented 89 and 95% relationship between polychaetes and environmental variables during first and second post monsoon season respectively (Fig. 7a & b). Organic carbon, salinity, sand and clay were the variables which affected polychaete abundance during first post monsoon and salinity, clay, OC and DO were the influencing parameters during second post monsoon season.

During pre-monsoon 95% relationship was observed between polychaetes and environmental variables (Fig.7c). Sediment characteristics (sand, silt, and clay), salinity and DO influenced macrobenthic polychaete abundance during this season. In the monsoon season 93% relationship was observed between polychaetes and environmental variables mainly DO, salinity and sediment characteristics (sand, silt and OC) (Fig. 7d). It was observed that preference of different species towards a particular parameter differed with the season (Figs.7a-d), as species in different groups are influenced by certain environmental parameter or sediment characteristics. Species located near the center of the ordination diagram indicated that these polychaetes are not influenced by any particular environmental variable (Figs.7a-d).

Canonical correspondence analysis (CCA) for non-polychaete groups and environmental variables

In the CCA biplot for non-polychaete macrobenthic groups, 96% of the relationship was observed between abundance of other macrobenthic groups and environmental variables during first post monsoon (Fig.8a), and it was 78% during second post monsoon (Fig.8b). DO, salinity, sand and OC were the important environmental variables influencing their abundance. Fish and oligochaetes preferred higher percentage of OC and low percentage of sand whereas, sipunculans preferred higher percentage of sand and lower OC (Fig.8a). Crustaceans preferred high OC whereas sipunculans preferred low OC (Fig.8b). During pre-monsoon, 79% of the relationship was observed between abundance and environmental variables (Fig.8c). OC, DO and salinity were the important environmental variables influencing abundance of other groups. During this season also crustaceans preferred higher OC. During monsoon the relationship between macrobenthos abundance and environmental variables was 91% (Fig.8d). Sediment characteristics (clay and silt) were the important environmental variables influencing abundance. In general, crustaceans preferred higher values of OC and clay and lower values of DO and salinity.

DISCUSSION

The present study was carried out in and around the Visakhapatnam Harbour selecting 24 stations which were topographically divided into two major areas - the Inner and the Outer Harbour. The seasonal variation in the abundance of macrobenthos in the harbour area is reported and discussed in relation to the influence of sediment characteristics and near bottom hydrological parameters.

Raman & Ganapati, (1986) conducted faunal studies in Visakhapatnam Harbour in 1976 and reported that polychaetes were the most diverse organisms in the macrobenthic community. In the present study 84 macrobenthic forms were reported, out of which 60 were polychaetes and 24 forms belonging other groups (non-polychaete). An earlier study of Visakhapatnam Harbour reported 38 polychaete species in northern arm in 1975 (Sarma & Ganapati, 1975), and a year later the number of polychaetes reported was reduced to 12 species (Raman & Ganapati, 1986). Raman, (1995) later reported higher abundance of opportunistic species in Visakhapatnam harbour including *Capitella capitata* and *Nereis glandicineta*. However, he reported 12 species when he carried out the study in 1976. The disappearance of species reported earlier was considered to be due to increase in the pollution during the intervening years (Raman, 1995). This indicates the number of polychaete species increased in the present study (38 in 1975 to 60 in 2008) to about 150% when compared to earlier studies (1975-76) in Visakhapatnam Harbour. Abundance of macrobenthic polychaetes, *Cossura coasta*, *Nephtys dibranchis*, *Cirratulus* sp., *Cossura* sp., *Prionospio* sp., *Poecilochaetus* sp., *Ancistrosyllis constricta* was high during this study. Among other macrobenthic groups, crustaceans (tanaids and amphipods) and sipunculans were the most abundant. Hence the possible reason for an increase in the number of polychaete species, is due to the increase in the nutrient concentration, and also the species which inhabited the semi-polluted conditions have been replaced by the pollution tolerant species or opportunistic species. The IH, which is relatively more polluted than outer harbour showed considerable higher nutrient concentrations arising due to untreated domestic sewage discharge and also low circulation of water with the open sea has promoted the rich biotic communities. The polychaete species abundance differed with Inner and Outer Harbour.

Inner Harbour (IH)

In IH, *Cirratulus* sp., *Nephtys dibranchis* and *Poecilochaetus* sp. were abundant followed by sipunculans (pea-nut worm) and tanaids (Table 3) which are carnivorous and preferred fine sediment, indicating fine sediment could sustain the carnivorous benthic organisms. Similar observations were made by Jayaraj *et al.* (2008a), in a tropical continental margin along the west coast of India, where the sediment was sandy silt. Sipunculans which showed dominance in IH and can live under normal

oceanic salinity (lower limit 18 to 26psu) and far less sensitive even if it increases up to 35psu. They are also capable of surviving on the sparse organic matter available on the upper layer of sediments (Murina, 1984). Raman and Ganapati, (1986) reported that members of the Cossuridae were more conspicuous in the transition zone (semi-polluted) between the IH and open sea. Amphipods were abundant in IH with silty substratum which may be preferred by amphipods that inhabit in mud or among detritus and this might also enable these organisms to feed on the organic matter present within the fine sediment. Generally, deposit feeder *Cirratulus* sp. was abundant followed by *Cossura coasta* during pre-monsoon season when the percentage of silt was high. From the properties of sediment dynamics it has been suggested that high silt-clay fraction in sediment contains more food particles (Sanders, 1958, 1960; Sanders et al., 1962; Jayaraj *et al.*, 2008b), and this may be the reason for the occurrence of more number of deposit feeders in high silt conditions. Generally, tanaids (crustacean) were more during the monsoon season followed by polychaetes, *Cirratulus* sp. and *Nephtys dibranchis* in IH. Crustaceans which also contributed to the overall macrobenthic abundance were commonly encountered in IH during all seasons except first post monsoon season, where sipunculans showed dominance over the crustaceans. CCA biplot for other macrobenthos and environmental variables supported the observation that sipunculans preferred higher percentage of sand and lower percentage of OC during first post monsoon season. During first post monsoon season, *Nephtys dibranchis* was abundant in IH (sand+silt). The high organic carbon at Port dry dock (IH-3) and low at Turning basin (IH-4) in IH showed low species diversity and can be concluded that both higher and lower organic carbon affects species diversity. Harkantra, (1982) and Harkantra & Parulekar, (1982) stated that ample or median range of organic carbon content supports the rich faunal density, however, low and high values of organic content shows poor fauna. In IH, Port dry dock (IH-3) and Turning basin (IH-4) were silty and sandy respectively. Silt content or increased percentage of fine sediment may help in retaining organic matter and this may be the reason why Port dry dock (IH-3) and Container berth-1 (OH-6) showed higher values of organic carbon which supported abundance of *Cossura coasta*.

During second post monsoon season, amphipods (crustaceans) which are mostly referred as carnivores were abundant in IH followed by the polychaete *Cirratulus* sp. Organic carbon was also high at most of the IH stations with silty substratum and higher abundance of deposit feeders *Cirratulus* sp. and *Poecilochaetus* sp. which is considered as an indicator of healthy bottom conditions. Dean, (2008) made similar observation where he found polychaete *Poecilochaetus serpen* common in nearby non-polluted areas of Barcelona.

Nutrients showed a unique trend in the concentration at Inner and Outer Harbour. During the entire season nitrate, nitrite, silicate and phosphate values were high in Inner Harbour when compared to Outer Harbour, may be due to the release of domestic sewage along with industrial wastes into the IH area. The nutrients in seawater will influence the phytoplankton community and abundance and this in turn will affect the benthic organisms which feed on phytoplankton. Mathew and Govindan, (1995) observed high concentration of nitrite in the inner creek of Mumbai and suggested such conditions are unhealthy for benthic communities. Ganapati and Raman, (1973) also reported the presence of heavy metallic ions, such as copper, iron and fluorides, phosphorus and oil from the industrial effluents into the Visakhapatnam harbour. This may be one of the reasons for low macrobenthic polychaete abundance in IH compared to OH.

Outer Harbour (OH)

In Outer Harbour, *Cossura coasta* was abundant followed by *Ancistrosyllis constricta*. *Cossura coasta* was the dominant species during the second post monsoon in Outer Harbour (silty sand). It is reported that species belonging to *Nephtys* and *Cossura* were occurred in sandy and silty habitat respectively (Jayaraj *et al.*, 2008a; Flint and Rabalais, 1980). During monsoon the sediment texture was mixed type (more or less equal proportion of sand, silt and clay) and tanaids were abundant indicating that tanaids are not substratum specific unlike amphipods. In OH crustaceans showed dominance only during second post monsoon and pre-monsoon seasons which is supported by CCA biplot results indicating preference of crustaceans to high percentage of OC during these seasons.

Sediment parameters like grain size, organic content and food availability are among the important factors which have been related to benthic community structure (Sanders, 1958; Kari, 2002). Abundance of *Cossura coasta* at Container berth-1 (OH-6) in OH indicated it as an organically polluted area. Polychaetes such as *Cossura coasta*, *Cirratulus* sp., and *Prionospio* sp. (deposit feeders) and *Nephtys dibranchis* (a carnivore) were abundant in OH suggesting different forms of polychaetes can reside in the OH indicating its semi-polluted conditions.

Comparative account on inner and outer harbour with respect macrobenthos species diversity

Outer Harbour showed higher species diversity compared to IH. Higher silt and sand composition was observed during pre-monsoon in both Inner and Outer Harbour. *Cossura coasta* and *Cirratulus* sp. were abundant in OH. Mixed type sediment texture (sand, silt and clay) was observed during monsoon season in both Inner and Outer Harbour with wide ranging organic carbon. During this season tanaids contributed nearly 30% to the total macrobenthos in IH, whereas in OH their contribution was negligible.

Temperature was high by 2-3 °C during pre-monsoon and monsoon season compared to post monsoon during which high abundance of *Cirratulus* sp. was observed. Mistri *et al.*, (2002) reported species tolerant to broad ranges of environmental variables such as temperature and salinity can persist well in benthic habitat. Jayaraj *et al.*, (2007) in his study in the northwest Indian shelf stated that, silt, temperature, dissolved oxygen, sand, temperature-depth and temperature-DO were the most significant factors in predicting the density of polychaetes.

Nephtys dibranchis, a typical euryhaline form and also a stenohaline form such as *Cossura coasta* were the two most abundant polychaete species present at higher salinity ranges from 29.2 to 32psu. When the salinity was low (23.8 to 26psu) during second post monsoon, stenohaline forms such as *Cossura coasta*, *Cirratulus* sp. and amphipods were abundant. Thus *Cossura coasta* can tolerate a wide range in salinity and can be termed as euryhaline which might have adapted to such conditions over the years in this habitat. This indicates that salinity at the benthic-pelagic region plays an important role in the distribution and abundance of macrobenthos. Similar observations were made by Raman & Ganapati, (1983), that in OH stenohaline cossurid, *Cossura coasta*, and cirratulid, *Cirratulus* sp. was found in large numbers. During low saline conditions in monsoon because of considerable amount of fresh water enters the harbour through a monsoon fed-stream (RamaRaju *et al.*, 1990), euryhaline tanaids were the most abundant macrobenthos.

It has been suggested that higher organic carbon can cause a decline in species diversity, abundance and biomass, possibly due to oxygen depletion and buildup of toxic by-products such as ammonia and sulphide (Jorgensen, 1977; Revsbech & Jorgensen, 1986; Snelgrove & Butman, 1994; Hyland *et al.*, 2005) and also low oxygen or hypoxic conditions are an important limiting factors in species distribution (Rainer & Fitzhardinge, 1981). DO levels below 2.5 mg. L⁻¹ are not merely considered as hypoxic, however during first post monsoon season, in station East quay berth-9 (IH-13), DO was low (2.3 mg.L⁻¹), and organic carbon was >3% and this may be the reason for the absence of polychaetes at this station. The inner harbour stations, IH-5 (Hindustan Shipyard), IH-7 (Fertilizer Warf) and IH-12 (East Quay Berth) had low polychaete abundance. Stations IH-5 and IH-7 receive a large amount of industrial effluents, and also IH-7 being a fertilizer warf which can have fertilizers inoculating in the water and in the sediment during the cargo loading and unloading and will add to the increase in the concentration of inorganic nitrogen and phosphorus. This might also have implication to IH-5 which is near to IH-7. The station IH-12 had higher organic carbon and low DO and this station is situated at the inlet of untreated domestic sewage. Also IH-12 has higher clay content compared to other stations in the inner harbour, and this might have adverse impact on the feeding ecology of organisms and thereby reduce their survival and abundance. Overall IH has

higher concentration of nitrite, nitrate and phosphate indicating eutrophic conditions compared to OH. Similar was the situation at East quay berth-7 (IH-12) during second post monsoon. Decreasing concentration of oxygen diminishes both species richness and diversity and the species composition is largely determined by the tolerance to oxygen deficiency (Johansson, 1997; Flemer *et al.*, 1999; Wu, 2002). Reduced oxygen content of the water might have led to adverse biological effects on the soft bottom macrobenthic communities (Saiz-Salinas, 1997). The CCA results also indicated that salinity, DO and sediment characteristics are the most influential factors for macrobenthic polychaete abundance however this varied with the season.

In some stations (IH-10, IH-11, IH-12 and IH-13) dissolved oxygen was 1 mg. L⁻¹ and the corresponding organic carbon values were high, which supports the reason for the reduced number of polychaetes. Sanders *et al.* (1968) predicted low diversity in shallow areas due to the depletion of oxygen by high organic matter. Harkantra *et al.* (1982) reported a lower abundance of benthic polychaetes in areas where organic matter was more than 3% indicating their avoidance to high organic matter. CCA biplot for polychaetes during pre-monsoon season also supported this phenomenon indicating polychaetes preferred higher values of salinity and lower values of OC.

The species diversity also varied with seasons and with respect to their occurrence in IH and OH. Sandy substratum supported high species diversity, richness and evenness during first post monsoon season. Coarse sediment and medium sized grains support rich benthic fauna (Rodrigues *et al.*, 1982; Ingole *et al.*, 2002) and such faunal-sediment relationships have also been reported by Sanders (1958, 1962). Sukumaran & Devi, (2009) while working in Mumbai port indicated that Shannon diversity index of 2 and above can be considered as fairly healthy benthic ecosystem. In the present study Shannon diversity index values showed fairly good benthic environment at stations which had high species diversity. Among all the seasons second post monsoon season showed low species diversity. Salinity during this season was low and might be a reason for depletion of species diversity. During pre-monsoon higher species diversity was observed which was supported by sandy substratum and high salinity, temperature and DO. During monsoon, sandy substratum and optimum dissolved oxygen supported high species diversity.

The polychaete, *Cossura coasta* was the most abundant species which was followed by *Ancistrosyllis constricta* in the stations which are more or less sandy and silty with moderate organic carbon during post-monsoon I season, suggesting *Cossura coasta* may not be a substratum specific species and can also withstand diverse environmental conditions such as high organic matter and low DO. Other species belonging to Cossuridae (*Cossura* sp.) was also common and abundant species during post-

monsoon I and II with considerable contribution to the total abundance during pre-monsoon season. *Nephtys dibranchis* was abundant at stations which were more or less sandy. *Nephtys dibranchis* preferred mostly sandy substratum because of higher interstitial space (Jayaraj *et al.*, 2008b). Hoey *et al.* (2004) also reported the presence of *Nephtys dibranchis* in the medium grained sediment during pre-monsoon with *Nephtys cirrosa* community in Belgium shelf. During second post monsoon also *Cossura coasta*, *Cirratulus* sp. and *Nephtys dibranchis* were the common polychaetes. *Cirratulus* sp. was dominant in the stations with a silty substratum. During pre-monsoon *Cossura coasta* is common in both IH and OH, but dominant only in OH (Table 3). These stations were mostly silty with sand also contributing to some extent with uneven organic carbon and DO, indicating this species can withstand varying DO. Raman and Ganapati (1983) also reported *Cossura coasta* species in the sandy-clay substratum with wide range of DO.

Cirratulus sp. and *Cossura coasta* were the common and dominant species in the stations with sandy-silt substratum and moderate OC (<3%). Musale & Desai (2010) observed higher abundance of polychaetes along the west coast of India in sandy substratum and they opined that this may be due to higher interstitial space for polychaetes to harbour. However, Jayaraj *et al.* (2007) observed that, polychaete species of family Cirratulidae and Cossuridae has the ability to survive in adverse conditions suggesting that these two polychaetes are cosmopolitan and can survive in both healthy and unhealthy benthic conditions. During monsoon season DO ranged from 0.7 to 5.9 mg. L⁻¹, where *Cossura coasta*, *Cirratulus* sp., and *Ancistrosyllis constricta* were common. Pollution resistant polychaete, *Cirratulus* sp., was abundant at the stations with clayey and sandy substratum and high organic carbon. Polychaetes such as, *Dorvillea* sp., *Cirratulus* sp., *Serpula* sp., *Nereis* sp., *Cossura coasta*, *Ancistrosyllis* sp., and *Prinospio* sp. were also reported, however their abundance was low. The environmental parameters and sediment characteristics play an important role in the distribution and abundance of polychaetes, and the distribution and abundance varied in the Inner and Outer Harbour areas as Inner Harbour is more polluted and this was also indicated by the presence of pollution indicator species.

It can be concluded that the number of polychaetes in terms of species diversity inhabiting the Visakhapatnam Harbour has been increased over the last 20 years. As mentioned earlier the higher organic carbon and lower oxygen concentration cause a decline in the species diversity. Most of the polychaetes reported in the present study were observed in the stations which had varying levels of DO concentration and low organic carbon levels. Also the outer section of the harbour which is semi-polluted compared to Inner Harbour had higher species diversity. This indicates that over the years forms which can tolerate varying levels of these environmental conditions must have prevailed.

The larval forms of the benthic organisms are pelagic and can disperse through natural physical processes mainly currents and can travel long distance. This plausible reason cannot be ruled out while explaining the increased species diversity. Also bioinvasion of these organisms from other part of the world and port hopping of species within India might bring some of the polychaete larvae which can survive and proliferate in a port environment as Visakhapatnam receives ships ballast water from almost all part of the world and from all the ports within India. However this needs further validation.

ACKNOWLEDGMENTS

We are grateful to The Director, CSIR-National Institute of Oceanography, Goa for his support and encouragement. This work was carried out as a part of Ballast Water Management Program, India, funded by Directorate General of Shipping, Government of India. This is a NIO contribution #####

REFERENCES

- Bismillah S.** (1986) Ecology of macrobenthos in relation to pollution in the north arm in Visakhapatnam harbour. Thesis, Andhra University, Waltair pp. 59
- Buchanan J.B.** (1984) Sediment analysis. In Holme N.A., McIntyre AD (eds) Methods for the study of marine benthos. *Oxford: Blackwell Scientific Publications*, pp 41-645.
- Clarke K.R. and Gorley R.N.** (2001) *PRIMER v5: User manual/Tutorial* (p. 91). Plymouth: PRIMER-E Ltd.
- Day J.H.** (1967) A monograph on the polychaete of Southern Africa Part-I and II, *trustees of The British Museum* (Natural history). London.
- Dean H.K.** (2008) The use of polychaetes (Annelida) as indicator species of marine pollution: a review. *Revista de Biologia Tropical* 56(4), 11-38.
- Flemer D.A., Kruczynski W.L., Ruth B.F. and Bundrick C.M.** (1999) The relative influence of hypoxia, anoxia, and associated environmental factors as determinants of macrobenthic community structure in a Northern Gulf of Mexico estuary. *Journal of Aquatic Ecosystem Stress and Recovery* 6, 311-328.
- Flint W.P. and Rabalais N.N.** (1980) Polychaete Ecology and Niche Patterns: Texas continental shelf. *Marine Ecology Progress Series* 3, 193-202.
- Ganapati P.N.** (1969) Biology of pollution in Visakhapatnam harbour. *Marine Pollution Bulletin* 16, 1-3.
- Ganapati P.N. and Raman A.V.** (1973) Pollution in the Visakhapatnam harbour. *Current Science* 42, 490-492.

- Ganapati P.N. and Raman A.V.** (1979) Organic pollution and *Skeletonema* blooms in Visakhapatnam harbour. *Indian Journal of Marine Sciences* 8, 184-187.
- Gosner K.L.** (1971) *Guide to identification of marine and estuarine invertebrates XIX*: 693.
- Harkantra S.N.** (1982) Studies on sublittoral macrobenthic fauna of the inner Swansea bay. *Indian Journal of Marine Sciences* 11, 75-78.
- Harkantra S.N. and Parulekar A.H.** (1987) Benthos off Cochin, Southwest Coast of India. *Indian Journal of Marine Sciences* 16, 57-59.
- Harkantra S.N., Rodrigues C.L. and Parulekar A.H.** (1982) Macrobenthos off the shelf of North Eastern Bay of Bengal. *Indian Journal of Marine Sciences* 11, 115-121.
- Hoey G.V., Degraer S. and Vincx M.** (2004) Macro-benthic community structure of the soft bottom sediments at the Belgium continental shelf. *Estuarine Coastal and Shelf Science* 59, 599-613.
- Hyland J., Balthis L., Karakassis I., Magni P., Petrov A., Shine J., Vestergaard O. and Warwick R.M.** (2005) Organic carbon content of sediments as an indicator of stress in the marine benthos. *Marine Ecology Progress Series* 295, 91-103.
- Ingole B.S., Rodrigues N. and Ansari Z.A.** (2002) Macrobenthic communities of the coastal waters of Dabhol, west coast of India. *Indian Journal of Marine Sciences* 31(2), 93-99.
- Jayaraj K.A., Jayalakshmi K.V. and Saraladevi K.** (2007) Influence of environmental properties on macrobenthos in the northwest Indian shelf. *Environmental Monitoring and Assessment* 127, 459-475.
- Jayaraj K.A., Jacob J. and Kumar P.K.D.** (2008a) Infaunal macrobenthic community of soft bottom sediment in a tropical shelf. *Journal of Coastal Research* 24(3), 708-718.
- Jayaraj K.A., Sheeba P., Raichandran C., Arun P.K., Praseeda K.S., Nisha P.A. and Rasheed K.A.** (2008b) Response of infaunal macrobenthos to the sediment granulometry in a tropical continental margin-southwest coast of India. *Estuarine Coastal and Shelf Science* 77, 743-754.
- Johansson B.** (1997) Behavioural response to gradually declining oxygen concentration by Baltic sea macrobenthic crustaceans. *Marine Biology* 129, 71-78.
- Jørgensen B.B.** (1977) The sulfur cycle of a coastal marine sediment (Limfjorden, Denmark). *Limnology Oceanography* 22, 814-832.
- Kalavati C., Raman A.V., Vaidehi J. and Barati V.R.** (1997) Pollution effect on planktonic ciliates in Visakhapatnam harbour (inland). *Indian Journal of Marine Sciences* 15, 131-132.
- Kari E.** (2002) Soft sediment benthic biodiversity on the continental shelf in relation to environmental variability. *Marine Ecology Progress Series* 232, 15-27.
- Kovach W.** (1998) *Multi-Variate Statistical Package*. Ver. 3.01, Pentraeth.
- Mathew A. and Govindan K.** (1995) Macrobenthos in the nearshore coastal system of Bombay. *Proceedings of National Academic Science, India* 65(B) IV.

- Mistri M., Fano E.L., Ghion F. and Rossi R.** (2002) Disturbance and community pattern of polychaetes inhabiting Valle Magnavacca (Valli di Comacchio, Northern Adriatic Sea, Italy). *P.S.Z.N. Marine Ecology* 23(1), 31-49.
- Murina G.V.V.** (1984) Ecology of Sipuncula. *Marine Ecology Progress Series* 17, 1-7.
- Musale A.S. and Desai D.V.** (2010) Distribution and abundance of macrobenthic polychaetes along the South Indian coast. *Environmental Monitoring and Assessment* 178, 423-436
- Parsons T.R., Maita Y. and Lalli C.M.** (1984) *A manual of chemical and biological methods for seawater analysis*. Oxford: Pergamon Press
- Rainer S.F. and Fitzhardinge R.C.** (1981) Benthic communities in an estuary with periodic deoxygenation. *Australian Journal of Marine and Freshwater Resource* 32, 227-243.
- RamaRaju V.S., Sarma V.V., Raju G.R.K. and SubbaRao V.** (1990) Mixing in Visakhapatnam harbour and nutrient inputs to the nearshore waters, east coast of India. *Mahasagar-Bulletin of the National Institute of Oceanography* 23(1), 49-54.
- Raman A.V.** (1995) Pollution effect in Visakhapatnam harbour, India: An overview of 23 years of investigations and monitoring. *Helgol Meeresunters* 49(1-4), 633-645.
- Raman A.V. and Ganapati P.N.** (1986) Benthic polychaete macrofauna and pollution in Visakhapatnam harbour, India. In Thompson MF, Sarojini R. and Nagbhushanam R. (eds) *Indian Ocean- biology of benthic marine organisms*. Oxford & IBH publishing, New Delhi, pp 463-484.
- Raman A.V. and Ganapati P.N.** (1986) Eutrophication in Visakhapatnam harbour, east coast of India. *Indian Journal of Marine Sciences* 18, 33-36.
- Raman A.V. and Ganapati P.N.** (1983) Pollution effects on Ecobiology of Benthic polychaetes in Visakhapatnam harbour (Bay of Bengal). *Marine Pollution Bulletin* 14(2), 46-52.
- Rathod V., Bhat K.L., Sudhakar U., Sarma V.V. and Parulekar A.H.** (1995) Pollution impacts on chaetognaths of the Visakhapatnam harbour and neritic waters, east coast of India. *Journal of Marine Biological Association of India* 37, 185-190.
- Revsbech N.P. and Jørgensen B.B.** (1986) Microelectrodes: Their use in microbial ecology. *Advances in Microbial Ecology* 9, 293-352.
- Rodrigues C.L., Harkantra S.N. and Parulekar A.H.** (1982) Sub-littoral meiobenthos of the northeastern Bay of Bengal. *Indian Journal of Marine Sciences* 11, 239-242.
- Saiz-Salinas J.I.** (1997) Evaluation of adverse biological effects induced by pollution in the Bilbao estuary (Spain). *Environmental Pollution* 96, 351-359.
- Sanders H.L.** (1958) Benthic studies in Buzzards Bay, I. Animal sediment relationships. *Limnology and Oceanography* 3, 245-258.
- Sanders H.L.** (1960) Benthic studies in Buzzards Bay, III. The structure of the soft bottom community. *Limnology and Oceanography* 5, 138-153.

Sanders H.L. (1968) Marine benthic diversity: A comparative study. *American Naturalist* 102(925), 243-282.

Sanders H.L., Goudsmith E.L. and Hampson G.E. (1962) Animal-sediment relationship. *Limnology and Oceanography* 3, 245-258.

Sarma V.V., Raju G.R.K. and Bose Babu T. (1982) Pollution characteristics and water quality in Visakhapatnam harbour. *Mahasagar* 15, 15-22.

Sarma A.L.N. and Ganapati P.N. (1975) Meiofauna of the Visakhapatnam harbour in relation to pollution. *Bulletin of Department Marine Science, University of Cochin* 7, 243-255.

Snelgrove P.V.R. and Butman C.A. (1994) Animal sediment relationship revisited: Cause versus effect. *Oceanography and Marine Biology Annual Review* 32, 111-177.

Sukumaran S. and Devi K.S. (2009) Polychaete diversity and its relevance in the rapid environmental assessment of Mumbai Port. *Current Science* 97 (10), 1439-1444.

Suresh R.R.V., Patnaik K.V.K.R.K., Ramesh U.S. and Misra S.C. (2012) A study on the seasonal variability of nearshore waves off Visakhapatnam, east coast of India. *International Journal of Innovative Research & Development* 10, 373-389.

ter Braak C.J.F. (1995) Ordination. In Jongman R.H.G., ter Braak C.J.F. and Van Tongeren O.F.R. (eds) *Data Analysis in Community and Landscape Ecology*. Cambridge University Press. Cambridge, pp 91-173.

Tripathy S.C., Kusuma Kumari B.A.V.L., Sarma V.V. and Ramana Murthy T.V. (2005) Evaluation of trophic state and plankton abundance from the environmental parameters of Visakhapatnam harbour and near-shore waters, east coast of India. *Asian Journal of Microbiology Biotechnology & Environmental Sciences* 7(4), 831-838.

Wakeel S.K. and Riley J.P. (1956) Determination of organic carbon in marine muds. *Journal of Du. Conseil International Exploration* 22, 180-183.

Wu R.S.S. (2002) Hypoxia: from molecular responses to ecosystem responses. *Marine Pollution Bulletin* 45, 35-45.

Legend for Tables

Table 1. Number of species(S), number of specimens (N), Margalef species richness (d), Pielou's evenness (J') and Shannon index (H) of macrobenthic polychaetes during (a) Post-monsoon I (b) Post-monsoon II (c) Pre-monsoon (d) Monsoon in Visakhapatnam harbour

Table 2. Hydrological parameters (Temperature, Salinity, Dissolved Oxygen) and Nutrients (Nitrite, Nitrate, Phosphate and Silicate) at the sampling stations during four seasons. PM-I; Post monsoon I, PM-II; Post monsoon II, PreM – Pre monsoon, Mon - Monsoon

Table 3. Schematic presentation of the variations in physical parameters, sediment characteristics and dominant macrobenthos reported in the Visakhapatnam port during different seasons. Percentage values in the parenthesis in front of the macrobenthos indicate their contribution to total macrobenthic abundance.

Legend for figures

Figure 1. Map showing sampling stations in and around the Visakhapatnam harbour.

Figure 2. Sediment texture and organic carbon (percentage) at the sampling stations in and around the Visakhapatnam harbour during different seasons (a) Post-monsoon I (b) Post-monsoon II (c) Pre-monsoon (d) Monsoon.

Figure 3. Total macrobenthic abundance at the sampling stations in and around the Visakhapatnam harbour during different seasons (a) Post-monsoon I (b) Post-monsoon II (c) Pre-monsoon (d) Monsoon.

Figure 4. Total polychaete abundance in different sampling stations in and around the Visakhapatnam harbour during different seasons (a) Post-monsoon I (b) Post-monsoon II (c) Pre-monsoon (d) Monsoon (e) abundance of different polychaete species during different seasons.

Figure 5. Total abundance of other macrobenthic (non-polychaete) groups in different stations in and around the Visakhapatnam harbour during different seasons (a) Post-monsoon I (b) Post-monsoon II (c) Pre-monsoon (d) Monsoon.

Figure 6. Dendrogram for hierarchical clustering of macrobenthic polychaetes with Bray-Curtis similarity indices during different seasons (a) Post-monsoon I (b) Post-monsoon II (c) Pre-monsoon (d) Monsoon.

Figure 7. Canonical correspondence analysis showing polychaete abundance and their relationship to environmental variables during different seasons (a) Post-monsoon I (b) Post-monsoon II (c) Pre-monsoon (d) Monsoon.

Figure 8. Canonical correspondence analysis showing macrobenthic other groups abundance and their relationship to environmental variables during different seasons (a) Post-monsoon I (b) Post-monsoon II (c) Pre-monsoon (d) Monsoon.

Table 1. Number of species (S), Number of specimens (N), Margalef species richness (d), Pielou's evenness (J'), Shanon index (H), of macrobenthic polychaetes during (a) post monsoon-I, (b) post monsoon-II, (c) Pre-monsoon and (d) Monsoon seasons in Visakhapatnam harbour.

(a)

<i>Stations</i>	<i>S</i>	<i>N</i>	<i>d</i>	<i>J'</i>	<i>H(log_e)</i>
IH-1	14	946	1.90	0.77	2.02
IH-2	2	33	0.29	1.00	0.69
IH-3	2	258	0.18	0.46	0.32
IH-4	2	367	0.17	0.88	0.61
IH-6	7	525	0.96	0.72	1.40
IH-7	1	33	0.00		0.00
IH-8	3	158	0.40	0.21	0.24
IH-9	1	50	0.00		0.00
IH-10	4	175	0.58	0.75	1.05
IH-11	1	25	0.00		0.00
OH-1	7	250	1.09	0.81	1.57
OH-2	11	933	1.46	0.60	1.44
OH-3	11	599	1.56	0.58	1.40
OH-4	3	308	0.35	0.23	0.25
OH-5	17	692	2.45	0.71	2.01
OH-6	8	759	1.06	0.43	0.89
OH-7	8	674	1.07	0.41	0.86
OH-8	4	108	0.64	0.88	1.22
OH-10	7	863	0.89	0.81	1.58
OH-11	8	567	1.10	0.67	1.40

(b)

<i>Stations</i>	<i>S</i>	<i>N</i>	<i>d</i>	<i>J'</i>	<i>H(log_e)</i>
IH-1	9	850	1.19	0.81	1.78
IH-2	10	725	1.37	0.82	1.88
IH-3	3	75	0.46	0.62	0.68
IH-4	4	825	0.45	0.79	1.09
IH-6	7	58	1.48	1.00	1.95
IH-8	5	151	0.80	0.59	0.95
IH-9	2	108	0.21	0.78	0.54
IH-10	1	8	0.00		0.00
IH-13	1	17	0.00		0.00
OH-1	9	508	1.28	0.75	1.65
OH-2	10	400	1.50	0.81	1.85
OH-3	8	458	1.14	0.39	0.81
OH-4	7	275	1.07	0.81	1.57
OH-5	6	451	0.82	0.39	0.70
OH-6	6	324	0.86	0.27	0.48
OH-7	7	318	1.04	0.57	1.11
OH-8	7	386	1.01	0.84	1.64
OH-10	10	401	1.50	0.55	1.28
OH-11	3	542	0.32	0.24	0.27

(c)

<i>Stations</i>	<i>S</i>	<i>N</i>	<i>d</i>	<i>J'</i>	<i>H(log_e)</i>
IH-1	16	2892	1.88	0.48	1.33
IH-2	8	375	1.18	0.83	1.73
IH-3	1	8	0.00		0.00
IH-4	3	83	0.45	0.99	1.09
IH-6	4	83	0.68	0.79	1.09
IH-9	2	75	0.23	0.99	0.69
IH-10	3	25	0.62	1.00	1.10
IH-13	1	8	0.00		0.00
OH-1	14	1283	1.82	0.66	1.74
OH-2	11	708	1.52	0.66	1.57
OH-3	4	108	0.64	0.81	1.12
OH-4	7	433	0.99	0.53	1.03
OH-5	13	467	1.95	0.79	2.04
OH-6	9	658	1.23	0.62	1.37
OH-7	11	496	1.61	0.46	1.10
OH-8	5	383	0.67	0.81	1.31
OH-10	6	142	1.01	0.75	1.34
OH-11	9	433	1.32	0.65	1.43


(d)

Stations	S	N	d	J'	H(log_e)
IH-1	17	974	2.33	0.66	1.87
IH-2	7	291	1.06	0.79	1.53
IH-3	15	1433	1.93	0.73	1.98
IH-4	4	318	0.52	0.65	0.91
IH-5	2	50	0.26	1.00	0.69
IH-6	7	147	1.20	0.82	1.60
IH-8	3	134	0.41	0.95	1.04
IH-9	2	75	0.23	0.77	0.54
IH-10	1	13	0.00		0.00
OH-1	9	173	1.55	0.83	1.82
OH-2	9	508	1.28	0.80	1.77
OH-3	9	182	1.54	0.73	1.60
OH-4	11	740	1.51	0.56	1.35
OH-5	7	249	1.09	0.88	1.71
OH-6	6	408	0.83	0.80	1.43
OH-7	9	658	1.23	0.50	1.11
OH-8	5	276	0.71	0.87	1.39
OH-9	3	158	0.40	0.93	1.02
OH-10	5	115	0.84	0.93	1.49
OH-11	4	191	0.57	0.74	1.03

Table 2. Hydrological parameters (Temperature, Salinity, Dissolved oxygen) and nutrients (Nitrite, Nitrate, Phosphate and Silicate) at different sampling stations during four seasons. PM-I; Post monsoon I, PM-II; Post monsoon II, PreM – Pre monsoon, Mon - Monsoon

Stations	Temperature ^o C				Salinity(psu)				D.O.(mg. L ⁻¹)				Nitrite(μM)			Nitrate(μM)			Phosphate(μM)			Silicate(μM)		
	PM-I	PM-II	PreM	Mon	PM-I	PM-II	PreM	Mon	PM-I	PM-II	PreM	Mon	PM-II	PreM	Mon	PM-II	PreM	Mon	PM-II	PreM	Mon	PM-II	PreM	Mon
IH-1	27	28	29	29	30	25.4	30.1	25	4.2	4.6	5.4	5.1	2.3	1.4	1.9	16.4	6	5.8	22	6.4	16	17	5.2	30
IH-2	27	28	29	30	29	24.9	31.1	25	4.9	4.5	5.4	4.5	3.9	1.3	3.8	6.1	2.7	3.8	65	4.7	47	27	6.7	23
IH-3	27	28	28	30	30	24	30.6	25	4.9	4.3	4.8	3.8	2.7	3.6	5	42	11	4.8	7.8	19	49	44	13	34
IH-4	27	28	28	30	30	25	31.6	24	4.9	4.9	4	4.5	1.9	1.5	1.7	35.3	10	16	23	10	27	24	11	200
IH-5	NA	28	28	29	NA	24.6	31.4	24	3.9	4.7	3.7	4.6	3.1	1.5	3.7	26.5	5.9	6.4	44	13	32	69	8.6	38
IH-6	26	27	28	29	30	24.7	31.4	27	6.2	5	5.4	5.9	0.8	0.6	0.6	0.9	2.7	0.6	8.4	1.9	3	11	9.1	17
IH-7	26	28	28	28	29	24.6	31.4	25	3.9	3.8	2.4	4.4	1.8	1.8	3	67.9	8.5	7.6	28	8.5	21	31	12	55
IH-8	27	28	31	29	29	25.1	30.6	24	4.4	3.6	2.1	3.8	2.7	2	6.5	39.1	15	2.5	27	19	50	88	24	81
IH-9	26	28	28	29	30	25	30.8	24	3.3	4.4	3.5	4.8	3.5	2.4	2.2	11.6	19	21	41	16	21	46	11	87
IH-10	27	28	28	29	29	26	30.7	23	3.1	4.1	1.1	3	0.9	1.1	5.8	27.5	20	1.3	19	16	46	45	24	117
IH-11	27	28	28	29	29	25.2	30.5	22	3.5	3	1	2.9	2.2	1.5	4.3	90.9	4.2	2.5	22	18	49	95	28	93
IH-12	27	28	28	29	29	25.2	31.1	21	3.1	2.9	1.1	0.7	1.4	1.3	4.6	132	13	1.2	26	18	108	139	31	103
IH-13	27	28	28	29	29	26	30.8	20	2.3	4.5	1.1	1	1.8	0.5	1.6	132	2.3	5	18	17	64	79	33	52
OH-1	26	27	29	29	30	23.8	30.9	25	5.9	7	6.1	5.4	0.2	1.3	1.2	2	3.1	3.6	1.2	4.4	8.7	2.6	18	36
OH-2	26	27	28	29	30	24.3	31.4	26	6.2	4.4	4.6	5.6	0.7	1.8	1.1	2.3	6.2	3.3	7.3	6.1	6.6	13	7.9	14
OH-3	26	27	28	29	30	24.2	31.5	26	6.1	5	5.1	5.4	0.2	0.8	1.4	1.3	2.5	5.5	1.6	3.3	11	9.5	7.9	14
OH-4	26	27	28	29	30	23.9	31.4	25	6.1	5.1	5	5.1	0.3	0.6	0.8	1.9	3	2.9	2.1	1.4	4.6	3.6	6.2	15
OH-5	26	27	28	29	30	24	31.6	26	6	4.4	5.8	5.8	0.2	0.5	1.1	0.9	2.3	1.8	1.5	1.7	9.6	11	3.9	14
OH-6	26	27	29	29	30	24	31.1	26	6.1	4.8	5.1	5.6	0.2	1.2	1.1	1.9	5.3	0.5	2.6	5.1	11	10	7.5	12
OH-7	26	27	28	29	30	24	31.6	26	6	5	4.6	5.6	0.3	11	0.7	0.7	3.7	2.6	2.6	4.2	5.3	10	19	13
OH-8	25	26	28	28	31	23.9	31.6	28	4.4	5.1	4.3	5.4	0.3	1	1	0.9	1.8	1.3	0.5	2	1.7	14	22	11
OH-9	27	28	28	29	30	24.8	31.2	25	4.4	5.2	4.6	4.6	1.6	1.8	2	39.8	5.8	6.1	26	9.6	21	28	5.8	33
OH-10	26	27	28	28	31	24.2	31.9	27	5.6	5.8	5.4	5.8	1.5	0.3	1.1	0.7	1.7	1.2	11	0.4	7.2	14	6.6	11
OH-11	26	27	28	30	30	24.1	31.4	28	6.2	6	5.1	4.6	0.8	0.9	0.9	0.9	2.7	2.7	5.9	3.6	7.3	11	7.1	4.2

Table 3. Schematic presentation of the variations in physical parameters, sediment characteristics and dominant macrobenthos reported in the Visakhapatnam port during different seasons. Percentage values in the parenthesis in front of the macrobenthos indicate their contribution to total macrobenthic abundance.



Inner harbour

Outer harbour

Inner harbour

Outer harbour

Tanaid (29.2%)

Cossura sp. (8.7 %)

Cirratulus sp. (13.4%)

Cossura coasta(18.7%)

Cirratulus sp. (8.9 %)

Cirratulus sp. (6.7%)

Poecilochaetus sp. (12.3%)

Cirratulus sp. (10 %)

Nephtys dibranchis (7.1%)

Cossura coasta (4.4 %)

Amphipod (8%)

Nephtys dibranchis (5.3 %)

-----Sediment

Sediment

Sand 45.67 (±20.91) %, Silt 33.05(±31.63)%,

Sand 44.39(±27.01)%, Silt 54.30(±26.64)%,

Clay 5.01 (±5.24)%, O.C.2.60(±1.43)%

Clay 1.31(±1.24)%, O.C. 2.37(±1.24)%

Sal= Salinity ; Temp=Temperature ; D.O.=Dissolved Oxygen, O.C.=Organic Carbon.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 4e




Figure 5




Figure 6




Figure 8