

## **Spatio-temporal variation of alkaline phosphatase activity in coastal waters off Trivandrum**

**Mamatha S. Shivaramu., Amruta K. Randive., Ritu Kumari, Manguesh Gauns,**

**LokaBharathi A. Ponnappakkam.,**

Biological Oceanography Division,  
CSIR-National Institute of Oceanography  
Dona Paula, Goa-403004, India

### **Corresponding author:**

\*Mamatha, S.S., Scientist  
Biological Oceanography Division,  
CSIR-National Institute of Oceanography  
Dona Paula, Goa-403004, India,  
E mail: mamatha@nio.org  
Ph no: +91 832 2450 254  
Fax: +91 832 2450 606

### **Summary**

Phosphatase is an extracellular enzyme which releases inorganic phosphate (Pi) from dissolved organic phosphate and indirectly organic carbon as nutrients for aquatic communities. Here, we have examined spatio-temporal variation in total alkaline phosphatase activity (APA) over a short period off Trivandrum, SW India. Sampling was at 50m water depth at 5, 15, 25 and 45 m for 5 consecutive days at 6hr intervals during post monsoon season. Total APA and phosphatase producing bacteria (PPB) were estimated along with pertinent environmental parameters. APA increased with depth upto  $3.98 \mu\text{M P hr}^{-1}$  at 45 m. Increase in pigment concentration with depth is responsible for increase in APA and Pi uptake. There is marginal increase in APA towards 18-24 hrs suggesting feeding activities of secondary producers. On the whole, chlorophyll and phaeophytin were responsible for nearly 45 and 55% variation in APA ( $p < 0.01$ ,  $p < 0.001$ ,  $n = 16$ ), respectively. Total bacterial count (TBC) was responsible for 32% ( $p < 0.05$ ,  $n = 16$ ) and total viable direct counts-aerobic (TVCa) for 24% ( $p < 0.05$ ,  $n = 16$ ) APA variation. About 38% ( $p < 0.01$ ,  $n = 20$ ) variation of APA was linked to chlorophyll at noon and 22% ( $p < 0.001$ ,  $n = 20$ ) to PPB at dawn. Thus, it is possible that bacteria and chlorophyll/phytoplankton could be responsible for variation in APA, with the latter contributing > the former at noon. Such studies would help to profile the fertility of coastal waters in terms of bioavailable Pi. Laboratory experiments are underway to help us discern the extent of light dependent contribution of chlorophyll/phytoplankton to APA and light independent participation of bacteria to the process.

**Key Words:** Phosphorus; Alkaline phosphatase activity; Chlorophyll; Phytoplankton; Bacteria

## **Introduction**

The availability of organic carbon and inorganic nutrients in water limits the biological productivity of aquatic environments. Of the essential elements involved in the biogeochemical cycle of the marine environment, phosphorus and nitrogen are most often considered as limiting factors. Dissolved and dead organic matter unusable by other organisms is subjected to microbial decomposition. Hence, the 'microbial loop' is important for recycling of phosphate and nitrogen in both water and sediment (Azam et al., 1983; 1995). The extensive number of microbial ectoenzyme activities detected in the water column has been noticed. Also the importance of heterotrophic bacteria and phytoplankton in bio-chemical/chemical processing and biogeochemical cycling in the ocean has been reported (Benitez-Nelson, 2000). Hydrolytic enzymes include alkaline phosphatase, chitinase, lipase, protease, aminopeptidase, glucosidase, and others. These enzymes convert organic matter to more labile form and thus influence the growth dynamics of different microbes and phytoplankton (Azam et al., 1995; Chrost, 1991; Martinez et al., 1996). Phytoplankton and bacteria can satisfy their phosphorus (P) requirements by producing hydrolytic enzymes such as alkaline phosphatase (AP) to access Pi from organic P. Pi gets limited in coastal areas with high nitrogen availability or in oceans where there is high rates of nitrogen fixation (Ammerman et al., 2003; Labry et al., 2005; Vidal et al., 2003; Yucel, 2018).

Alkaline phosphatases have a broad range of organic substrate specificities, and therefore hydrolyze a wide variety of organic phosphomonoesters (Ammerman, 1991). These enzymes are attached to cell surface or, are freely dissolved in the water column resulting from cell lysis or excretion (Hoppe, 2003; Jansson et al., 1988; Li et al., 1998). Upon enzyme hydrolysis, phosphomonoesters release inorganic P (Pi) into the water, along with their organic moiety, thereby increasing Pi availability for planktonic, as well as benthic organisms in shallow marine systems (Dyhrman and Ruttenberg, 2006; Koch et al., 2009). APA has been identified and associated with major groups of algae, cyanobacteria ((Dyhrman and Ruttenberg, 2006; Martinez et al., 1996) and heterotrophic bacteria (De Souza et al., 2000) and also studied in seawater (Koch et al., 2009; Mamatha et al., 2012) and sediment (Taga and Kobori, 1978). Generally, it is reported that APA gets expressed when P is limiting and suppressed when P level is replete. It is presumed that APA promotes organic P mineralization and recycling within P-limited systems (Ammerman et al., 2003; Koch et al., 2009; Labry et al., 2005; Mamatha et al., 2015; Sebastian et al., 2012; Vidal et al., 2003).

Concentration of P in coastal waters could be increased by either land run off or by upwelling process. In the rainy season (June-September), strong south west monsoonal winds cause coastal upwelling in Arabian Sea, which is one of the most productive areas in the world (Naqvi et al., 2010;

Prell et al., 1991). However, upwelling weakens in the southernmost part of the west coast of India by October, and as a result, the high concentrations of nutrients available at the surface water get exhausted due to autotrophic production (Peetersa et al., 2002). During the post monsoon season, the weaker north east monsoon winds result in deeply stratified water and deeper euphotic zone (Peetersa et al., 2002; Pillai et al., 2000). Based on this information, we have examined the spatio-temporal variation in APA along with other environmental parameters over short period off Trivandrum to appreciate the extent of changes in fertility as shown by the variation in APA. Previously, Mamatha et al. (2015) have noticed that low APA could be indicative of P sufficiency in coastal waters and higher activity suggestive of deficiency in off-shore waters off Trivandrum during upwelling season. In this communication on spatio-temporal variation during post upwelling, there is an increase in APA with depth due to increase in pigment and also Pi utilization without any temporal changes. The paper discusses these remarkable aspects.

The study station was off Trivandrum coast, south west of India (N 8° 26.04', E 76° 30') in the eastern Arabian Sea. Sea water samples were collected onboard FORV Sagar Sampada (cruise no. 282) during post monsoon season (November 2010) at lat 08°27.742'N and long: 76°46.709'E at maximum water depth of 50 m. The sampling was done at 4 depths i.e. 5, 15, 25 and 45 m bss (below sea surface) at 6 hourly intervals for 5 days. The sampling hours were 0 (24), 6, 12 and 18 hrs where 6 and 18 hrs were designated as dawn and dusk and 12 and 24 hrs as noon and midnight, respectively. The data with time intervals 6, 12, 18 and 24 hrs for 5 days were averaged to discern the spatio-temporal variation (supplementary table S1). Each data point is derived from triplicates.

Samples for dissolved oxygen (DO) were collected in 125 mL stoppered glass bottles avoiding air bubbles and were immediately fixed with Winkler's reagents. Fixed samples were stored in the dark until analysis (Strickland and Parsons, 1965). Sub-samples were taken for analysis of various parameters like total APA, chlorophyll a, phaeopigment, phosphate and other nutrients. Seawater sub-samples for total bacterial count (TBC) were collected in sterile containers immediately after retrieval and fixed with buffered formalin. Seawater sub-samples for other microbiological analyses were similarly collected and stored at 4 °C till analysis.

Salinity and temperature were derived from the Sea Bird CTD connected to the rosette Niskin sampler that operated vertically all the time. The pH was checked onboard using pH meter.

Winkler's titrimetric method (Strickland and Parsons, 1965) was used for estimating the dissolved oxygen (DO) concentration onboard using a Dosimeter (Metrohm 785 DMP Titrimo). Nutrients were also analyzed on board by SKALAR auto analyzer as described by Wurl (2009).

Chlorophyll *a* (Chl*a*) and phaeophytin (phaeo) concentrations were determined fluorometrically (Turner Designs, USA) by filtering 1 L water samples from each depth with GF/F filters and extracting with 10 mL of 90% acetone in the dark for 24 hrs under refrigeration. The extract was measured following UNESCO (1994) protocol. The readings were measured before and after acidification using Turner Design (10AU) fluorometer and then both Chl*a* and phaeo concentrations were calculated.

Total bacterial abundance (TBC) was counted using epifluorescence microscopy (Olympus Fluorescent Microscope Model BX61) by acridine orange direct count (AODC) method (Hobbie et al., 1977) and expressed as numbers per litre (nos L<sup>-1</sup>). Total direct viable counts (TVC), were estimated as outlined by Kogure et al. (1980) using a mixture of piromedic, pipemedic, nalidixic acid and yeast extract.

The end point dilution method (MPN=Most Probable Number) is based on a series of dilutions prepared from a sample, where selected liquid media are inoculated with each dilution using 3 parallels. After the incubation period, estimation of microbial count was done, using statistical tables based on the number of positive wells (showing colour formation i.e. colourless to yellow colour). Three tube method was suitably adopted for microtiter plate method. Phosphatase producing bacteria (PPB) were enumerated by most probable numbers (PPB-MPN) by microtitre plate method using para-nitrophenol phosphate (pNPP) as substrate (1 mM) (Fig 1s). The hydrolysis of phosphate was indicated by the release of colored para-nitrophenol (pNP) from colourless paranitrophenol phosphate (pNPP) in proportion to the Pi released. The samples were incubated in the dark at room temperature (28±2°C) for scoring positive and negative growth/reactions. Scoring/calculations were done using McCrady's table (Rodina, 1972). It is assumed that the color development is attributed only to bacteria and free enzymes in the diluted inoculum.

The alkaline phosphatase activity (APA, EC 3.1.3.1) was estimated using the method of Taga and Kobori (1978) and Mamatha et al. (2015). Measurements were run against a control containing an equivalent volume of the seawater samples previously autoclaved. The results represent the average of triplicates corrected for controls. Experimental values were calculated against a standard (para nitro phenol, pNP) curve prepared to cover concentrations from 10 nM to 1000 nM.

Microbiological data was log(x+1) transformed for statistical analysis. Inter-relationships between biological and environmental parameters were examined using Statistica ver. 6.0 and analysis tool pack in Microsoft Excel. Analysis of variance (ANOVA) was carried out to find out significant spatio-temporal variations in pertinent parameters.

In the present study, there is little variation in salinity with minimum of 33.07 and maximum of 35.46. The increase of salinity from surface to bottom is evident but it is not statistically significant. Variation in temperature and pH were also minimal as shown in the table 1. The dissolved oxygen in the water column ranged between 1.19 ml L<sup>-1</sup> in 45 m depth to 4.42 ml L<sup>-1</sup> in 5m layer (Table 1). There was significant variation in vertical distribution with depth and time in temperature and DO ( $p < 0.05$ ). Higher concentration of nitrate (20.02  $\mu\text{M}$ ) was measured at 45m while nitrite (1.55  $\mu\text{M}$ ) was measured at surface waters (Table1). Nitrate and nitrite showed significant variation with depth but not with time. The average concentration of phosphate over the study period was  $0.268 \pm 0.246$   $\mu\text{M}$  ( $n=80$ ) and it ranged from 0.08-2.37  $\mu\text{M}$ . Phosphate concentration was higher at 25m depth (2.37  $\mu\text{M}$ ) than at the 5m depth (0.08  $\mu\text{M}$ ). However, the spatial and temporal variation was not statistically significant.

The chlorophyll *a* and bacterial numbers can effectively represent algal and bacterial biomass. Chl*a* concentration varied from 0.71-2.90  $\mu\text{g L}^{-1}$  with an average  $1.39 \pm 0.50$   $\mu\text{g L}^{-1}$  (Table 1). Surprisingly, it was high at 45 m depth (2.9  $\mu\text{g L}^{-1}$ ) at all time intervals except noon when it is uniform throughout (Fig. 1a). Like chlorophyll, phaeopigment also follows the same trend in distribution (Fig. 1b).

TBC ranged from  $2.4 \times 10^8$  to  $1.54 \times 10^{10}$  cells L<sup>-1</sup> ( $5 \times 10^9 \pm 3 \times 10^9$  cells L<sup>-1</sup>) with maximum at the 45 m waters showing one order higher than the other depths (Table 1). TVCa was one order less than the TBC and was marginally higher at 45m where it varied from  $6.37 \times 10^8$ - $8.16 \times 10^9$  cells L<sup>-1</sup> (av.  $3.7 \times 10^9 \pm 2 \times 10^9$  cells L<sup>-1</sup>). PPB-MPN ranged from ndl- $1.5 \times 10^6$  cells L<sup>-1</sup> (av.  $0.27 \times 10^6 \pm 0.46 \times 10^6$  cells L<sup>-1</sup>). Though there are large standard deviations in the average values of the bacterial parameters, ANOVA showed that there was no significant variation in distribution with depth and time (Fig. 2).

Total APA of the whole water column ranged from ndl to 13.79  $\mu\text{M P hr}^{-1}$  ( $n=80$ ) (Table 1). It increased with depth to a maximum of 3.98  $\mu\text{M P hr}^{-1}$  ( $n=16$ , Fig. 3) at 45 m ( $p < 0.05$ ) corresponding to increase in chlorophyll and phaeophytin concentration. There is not much variation with time, however, marginal increase towards 18 to 24 hrs was discernible. Two way ANOVA showed that depth variation is stronger ( $p < 0.05$ ) than the temporal variation in APA.

On the whole, APA is positively related to depth ( $r=0.63$ ,  $p < 0.05$ ,  $n=80$ ), and negatively with DO ( $r=-0.594$ ,  $P < 0.05$ ,  $n=16$ ). Nitrate influenced 41% ( $r=0.637$ ,  $p < 0.01$ ) variation in APA. Chlorophyll and phaeophytin were responsible for 45 and 55% variation in APA, respectively ( $p < 0.05$ ,  $p < 0.001$ ,  $n=16$ ) (Table 2). Bacterial fractions examined contributed to the changes in APA with the TBC being responsible for nearly 32% ( $p < 0.05$ ,  $n=16$ ) and TVCa for 24% ( $p < 0.05$ ,  $n=16$ ). Time dependent

study showed that 38% ( $p < 0.01$ ,  $n = 20$ ) variation of APA at noon was linked to chlorophyll and 22% ( $p < 0.001$ ,  $n = 20$ ) variation in APA to PPB-MPN at dawn (Table 2).

The spatial variation (depth) is discernible in these waters (Nyadjro et al., 2012). The variation with temperature and DO confirms that there is stratification in these shallow coastal waters during this season but restricted to physico-chemical parameters. The higher concentration of nitrate at the bottom suggests that the source could be from sediment. Chl $a$  and phaeophytin were also high at 45 m all the time, suggesting that the settling of the pigments was high throughout the observation. The chlorophyll maxima was observed at noon at all the depths. This peaking could be a reflection of the penetration of PAR (photosynthetically active radiation) upto 45m depth. However, the depth of the euphotic zone could vary in different waters. Qasim (1982) reported that euphotic zone in the southern Arabian Sea was 60 m and in the northern part it was 40 m. Pillai et al. (2000) also found that the average depth of the euphotic zone in the west coast is 60 m during post monsoon season. Lack of significant variation in the different bacterial parameters with depth and time suggests that this could be due to their stronger association with phytoplankton (Reimann et al., 1999; Rooney-Varga et al., 2005). Rise in bacterial population at the bottom may be a consequence of nocturnal feeding and excretion by benthic organisms. Nandakumar and Damodaran (1998) found that there is a diurnal variation in the feeding intensity of speckled shrimps at Kochi. They feed more in the night than day. This benthic activity may enrich the bottom area with essential nutrients subsequently allowing bacteria including PPB to proliferate rapidly. Thus there is less heterogeneity in the microbial distribution in the coastal waters, suggesting that the representative sampling can be at any depth.

In the present study, APA peaks were observed from dusk to dawn (1/4 observations-dusk, 1/4-midnight and 2/4 dawn) (Fig.3). These APA peaks could be linked to prolonged grazing by zooplankton. Their sloppy feeding generally leads to release of phytoplankton exudates (Fouilland et al., 2013; Lignell, 1990; Yucel, 2018). Besides, higher APA at 45m layer could also be a reflection of higher phosphatase producing bacterial population from sediments below (Ayyakkannu and Chandramohan, 1970; 1971; Barik et al., 2001). Increased expression of APA with depth is noticeable in spite of same level of Pi. Hence, it is inferred that uptake rate could also be high with depth. On the other hand, Kobori and Taga (1979) found that APA was high at the surface layer and decreased with increasing depth at Tokyo Bay. This difference could be because the water is shallower in the present study than Tokyo Bay. Experimentally, Lin et al. (2011, 2012) have shown that bacterial contribution to APA could also be as important as phytoplankton. This inference could be particularly true for coastal waters.

The relation of APA with DO and nitrate are indirectly related to depth. APA relates with both pigments and bacterial parameters. The variation of *Chla* and phae with nitrate could be partly due to their requirement of this nutrient and also because nitrate and *Chla* co-vary with depth. The relationships between APA and biological parameters are discernible in scatter plots (Fig S2). Thus, it is possible that both bacteria ( $r=0.561$ ,  $p<0.05$ ) and *Chla* ( $r=0.670$ ,  $p<0.01$ ) could be responsible for variation in APA, with *Chla* ( $r=0.619$ ,  $p<0.01$ ) contribution being prominent at noon. Kwon et al. (2011) in the northern part of Gamak Bay, Korea and Hino (1988) in Lake Barato, Japan also found strong linear relation between APA and *Chla*. Thus, it could be inferred that major part of APA might have been induced by phytoplankton during the day time. Chrost and Overbeck (1987) also observed that phytoplankton were major APA producers in the photic zone of the lake plu\see of East Holstain (North Germany). During summer or winter, phytoplankton APA constituted, on an average, 49% of the total APA in the plu\see water but bacteria were the dominant APA producers in winter (41.3-44.9%). Increased phaeopigment is indicative of degradation of *Chla*. When *Chla* degrades, dissolved phosphate gets released into the environment and may indirectly supply the Pi requirement to phytoplankton and microbial communities. Dissolved AP can also be liberated into the environment through the lysis of dead phytoplankton cells (Chrost et al., 1991). Taga and Kobori (1978) found positive relationship between phosphatase and Pi in the eutrophic sea water samples in Tokyo Bay. In our study, the phosphate concentration was similar over time and depth (ca 0.2  $\mu\text{M}$ ). This lack of variation could be attributed to the balance between uptake and release of this nutrient by the water column community. Induction of APA by phytoplankton may be ecologically significant, allowing dominance by these organisms under phosphate limiting conditions (Kwon et al., 2011; Ly et al., 2014). Thus, the synchronization between the APA- *Chla* and APA-TBC ( $n=16$ ) during whole time series suggests that both bacteria and phytoplankton contribute to total APA.

We therefore conclude that there is a positive synchronization of APA with biotic variables like *Chla*, phaeophytin and bacterial abundance. Phytoplankton contribution to variation in total APA is more in noon while bacteria contributes during dawn ( $r=0.465$ ,  $p<0.001$ ). Like phytoplankton, bacteria also have a major role in continuously making this important nutrient, phosphate, bioavailable to the surrounding waters especially in the sub euphotic layers. Continuing the observation for longer time will strengthen our inference about the extent of contribution by primary producers and bacteria to APA in coastal system. Such studies will help us to understand the changes in productivity of coastal waters in space and time.

The authors are grateful to Directors, CSIR-National Institute of Oceanography for the facilities provided. They also express their gratitude to Ministry of Earth Science and Director, CMLRE,

Kochi, India for providing FORV Sagar Sampada for the above study. The authors also thank all the participants, Captain and crew members of Sagar Sampada cruise no. 282 for their constant support during sample collection and analyses onboard. This work was carried out under “Supra institutional (SIP 1301)” and “PSC 0206” projects of Council of Scientific and Industrial Research, India. This is NIO contribution no xxxx.

## References

- Ammerman, J.W., 1991. Role of ecto-phosphohydrolases in phosphorus regeneration in estuarine and coastal ecosystems. in: Chrost, R.J. (ed.), *Microbial enzymes in aquatic environments*. Springer-Verlag, New York, pp. 165-186.<https://doi.org/10.1007/978.1.4612.3090.8.10>.
- Ammerman, J.W., Hood, R.R., Case, D.A., Cotner, J.B., 2003. Phosphorus deficiency in the Atlantic: An emerging paradigm in oceanography. *Eos, Transactions American Geophysical Union*, 84(18), 165-170.<https://doi.org/10.1029.2003/EO180001>.
- Ayyakkannu, K., Chandramohan, D., 1970. On the occurrence and distribution of phosphobacteria in the marine environment at Porto-Novo. *Curr.Sci.* 39, 398-399
- Ayyakkannu, K., Chandramohan, D., 1971. Occurrence and distribution of phosphate solubilizing bacteria and phosphatase in marine sediments at Porto Novo. *Mar. Biol.* 11(3), 201-205.<https://doi.org/10.1007/BF00401268>.
- Azam, F., Fenchel, T., Field, J.G., Gray, J.S., Meyer-Reil, L.A., Thingstad, F., 1983. The ecological role of water-column microbes in the sea. *Mar. Ecol. Prog. Ser.* 10, 257–263. <https://doi.org/10.3354/meps010257>.
- Azam, F., Smith, D.C., Long, R.A., Steward, G.F., 1995. Bacteria in oceanic carbon cycling as a molecular problem. In: Joint, I. (eds.), *Molecular Ecology of Aquatic Microbes*, NATO ASI Series,38, Springer-Verlag, Berlin,pp. 39–54.<https://doi.org/10.1007/978.3.642.79923.5.3>.
- Barik, S.K., Purushothaman, C.S., Mohanty, A.N., 2001. Phosphatase activity with reference to bacteria and phosphorus in tropical freshwater aquaculture pond systems. *Aquacult. Res.* 32, 819-832.<https://doi.org/10.1046/j.1355-557x.2001.00619.x>.
- Benitez-Nelson, C.R., 2000. The biogeochemical cycling of phosphorus in marine systems. *Earth Sci. Rev.* 51, 109-135.[https://doi.org/10.1016/S0012-8252\(00\)00018-0](https://doi.org/10.1016/S0012-8252(00)00018-0).
- Chrost, R.J., Overbeck, J., 1987. Kinetics of alkaline phosphatase activity and phosphorus availability for phytoplankton and bacterioplankton in lake plu\see (North German Eutrophic Lake). *Microb. Ecol.* 13 (3), 229-248. <https://doi.org/10.1007/BF02025000>.
- Chrost, R.J., 1991. Environmental control of the synthesis and activity of aquatic microbial ectoenzymes. in: Chrost, R.J. (ed.), *Microbial enzymes in aquatic environments*. Springer-Verlag, New York Inc.,pp. 29-59.[https://doi.org/10.1007/978-1-4612-3090-8\\_3](https://doi.org/10.1007/978-1-4612-3090-8_3)
- De Souza, M.J.B.D., Nair, S., Chandramohan, D., 2000. Phosphate solubilizing bacteria around the Indian Peninsula. *Indian J. Mar. Sci.* 29, 48-51.



- Dyhrman, S.T., Ammerman, W.J., Van Mooy, B.A.S., 2006. Microbes and the marine phosphorus cycle, microbes and major elemental cycles. *Oceanogr.* 20(2), 196-199.<https://doi.org/10.5670/oceanog.2007.54>.
- Dyhrman, S.T., Ruttenberg, K.C., 2006. Presence and regulation of alkaline phosphatase activity in eukaryotic phytoplankton from the coastal ocean: Implications for dissolved organic phosphorus remineralization. *Limnol. Oceanogr.* 51(3), 1381-1390.<https://doi.org/10.4319/lo.2006.51.3.1381>.
- Fouilland, E., Tolosa, I., Bonnet, D., Bouvier, C., et al., 2014. Bacterial carbon dependence on freshly produced phytoplankton exudates under different nutrient availability and grazing pressure conditions in coastal marine waters. *FEMS Microbiol.Ecol.* 87(3),757-769.  
<https://doi.org/10.1111/1574-6941.12262>.
- Hino, S., 1988. Fluctuation of algal alkaline phosphatase activity and the possible mechanisms of hydrolysis of dissolved organic phosphorus in Lake Barato. *Hydrobiol.* 157(1), 77-84.<https://doi.org/10.1007/BF00008812>.
- Hobbie, J.E., Daley, R.J., Jasper, S., 1977. Use of nuclepore filters for counting bacteria by fluorescence microscopy. *Appl. Environ. Microbiol.* 33(5), 1225-1228.
- Hoppe, H.G., 2003. Phosphatase activity in the sea. *Hydrobiol.* 493(1/3), 187-200.<https://doi.org/10.1023/A:1025453918247>.
- Jansson, M., Olsson, H., Pettersson, K., 1988. Phosphatases; origin, characteristics and function in lakes. *Hydrobiol.* 170(1), 157-175.<https://doi.org/10.1007/BF00024903>.
- Kobori, H., Taga, N., 1979. Phosphatase activity and its role in the mineralization of organic phosphorus in coastal sea water. *J. Exp. Mar. Biol. Ecol.* 36(1), 23-39.[https://doi.org/10.1016/0022-0981\(79\)90098-4](https://doi.org/10.1016/0022-0981(79)90098-4).
- Kogure, K., Simidu, U., Taga, N., 1980. Distribution of viable marine bacteria in neritic seawater around Japan. *Canadian J. Microbiol.* 26(3), 318-323.<https://doi.org/10.1139/m80.052>.
- Koch, M.S., Kletou, D.C., Tursi, R., 2009. Alkaline phosphatase activity of water column fractions and seagrass in a tropical carbonate estuary, Florida Bay. *Estuarine, Coastal and Shelf Science.* 83, 403-413.<https://doi.org/10.1016/j.ecss.2009.04.007>.
- Kwon, H.K., Oh, S.J., Yang, H.S., 2011. Ecological significance of alkaline phosphatase activity and phosphatase-hydrolyzed phosphorus in the northern part of Gamak Bay, Korea. *Mar. Pollut. Bull.* 62(11), 2476-2482.<https://doi.org/10.1016/j.marpolbul.2011.07.027>.
- Labry, C., Delmas, D., Herbland, A., 2005. Phytoplankton and bacterial alkaline phosphatase activities in relation to phosphate and DOP availability within the Gironde plume waters. *J. Exp. Mar. Biol. Ecol.* 318, 213-225.<https://doi.org/10.1016/j.jembe.2004.12.017>.
- Li, H., Veldhuis, M.J.W., Post, A.F., 1998. Alkaline phosphatase activities among planktonic communities in the northern Red Sea. *Mar. Ecol. Prog. Ser.* 173, 107-115.
- Lignell, R., 1990. Excretion of organic carbon by phytoplankton: its relation to algal biomass, primary productivity and bacterial secondary productivity in the Baltic Sea. *Mar. Ecol. Prog. Ser.* 68(1-2), 85-99.<http://dx.doi.org/10.3354/meps068085>.

- Ly, S.J., Philippart, C.J.M., Kromkamp, J.C., 2014. Phosphorus limitation during a phytoplankton spring bloom in the western Dutch Wadden. *Quaternary Res.* 88,109-120.<http://dx.doi.org/10.1016/j.seares.2013.12.010>.
- Mamatha, S.S., Gobika, A., Janani, S., 2012. Phosphate solubilizing bacteria and alkaline phosphatase activity in coastal waters off Trivandrum. *J. Coast. Env.* 3(1), 89-100.
- Mamatha, S.S., Malik, A., Varik, S., Paravthi, V., Jineesh, V.K., Gauns, M.U., LokaBharathi, P.A.,2015. Alkaline phosphatase activity at the southwest coast of India: a comparison of locations differently affected by upwelling. *J. Sea Res.*95, 196-205. <https://doi.org/10.1016/j.seares.2014.06.002>.
- Martinez, J., Smith, D.C., Steward, G.F., Azam, F., 1996. Variability in ectohydrolytic enzyme activities of pelagic marine bacteria and its significance for substrate processing in the sea. *Aquat. Microb. Ecol.* 10(3),223–230.<https://doi.org/10.3354/ame010223>.
- Nandakumar, G., Damodaran, R., 1998. Food and feeding habits of the speckled shrimp *Metapenaeusmonoceros* (fabricius). *J. Mar. Biol. Ass. India*, 40 (1 & 2), 30-43.
- Naqvi, S. W. A., Moffett, J. W., Gauns, M. U., Narvekar, P. V., Pratihary, A. K., Naik, H., Shenoy, D. M., Jayakumar, D. A., Goepfert, T. J., Patra, P. K., Al-Azri, A., Ahmed, S. I., 2010. The Arabian Sea as a high-nutrient, low-chlorophyll region duringthe late Southwest Monsoon. *Biogeosciences*,7, 2091-2100.<https://doi.org/10.5194/bg.7.2091.2010>.
- Nyadjro, E.S., Subrahmanyam, B.,Murty,V. S. N., Shriver, J. F., 2012. The role of salinity on the dynamics of the Arabian Sea mini warm pool. *J. Geophys. Res.* 117, C09002, 1-12.<https://doi.org/10.1029/2012.JC.007978>.
- Citation: Nyadjro, E. S., B. Subrahmanyam, V. S. N. Murty, and J. F. Shriver (2012), The role of salinity on the dynamics of the Arabian Sea mini warm pool, *J. Geophys. Res.*, 117, C09002, doi:10.1029/2012JC007978.
- Citation: Nyadjro, E. S., B. Subrahmanyam, V. S. N. Murty, and J. F. Shriver (2012), The role of salinity on the dynamics of the Arabian Sea mini warm pool, *J. Geophys. Res.*, 117, C09002, doi:10.1029/2012JC007978.
- itation: Nyadjro, E. S., B. Subrahmanyam, V. S. N. Murty, and J. F. Shriver (2012), The role of salinity on the dynamics of the Arabian Sea mini warm pool, *J. Geophys. Res.*, 117, C09002, doi:10.1029/2012JC007978.
- Peetersa, F.J.C., Brummera, G.J.A., Ganssen, G., 2002. The effect of upwelling on the distribution and stable isotope composition of *Globigerina bulloides* and *Globigerinoidesruber* (planktic foraminifera) in modern surface waters of the NW Arabian Sea. *Global and Planetary Change*, 34(3), 269-291.[https://doi.org/10.1016/S0921-8181\(02\)00120-0](https://doi.org/10.1016/S0921-8181(02)00120-0).
- Pillai, V.N., Pillai, V.K., Gopinathan C.P., Nandakumar, A., 2000. Seasonal variations in the physico-chemical and biological characteristics of the eastern Arabian Sea. *J. Mar. Biol. Ass. India*, 42 (1&2), 1-20.

Prell, W.L., Niitsuma, N., et al., 1991. Proceedings of the Ocean Drilling Program, Scientific Results, 117, 257-267.

Qasim, S.Z., 1982. Oceanography of the northern Arabian Sea. Deep Sea Res. Part A, 29(9), 1041-1068.[https://doi.org/10.1016/0198-0149\(82\)90027-9](https://doi.org/10.1016/0198-0149(82)90027-9).

Riemann, L., Steward, G.F., Fandino, L.B., Campbell, L., Landry, M.R., Azam, F., 1999. Bacterial community composition during two consecutive NE monsoon periods in the Arabian Sea studied by denaturing gradient gel electrophoresis (DGGE) of rRNA genes. Deep Sea Res. II 46 (8-9), 1791-1811.[https://doi.org/10.1016/S0967-0645\(99\)00044-2](https://doi.org/10.1016/S0967-0645(99)00044-2).

Rodina, A.G., 1972. Methods in Aquatics Microbiology. Colwell, R.R., Zambruski, M.S. (Eds) Baltimore-London-Tokyo, University Park Press, pp.461.

Rooney-Varga, J.N., Giewat, M.W., Savin, M.C., Sood, S., LeGresley, M., Martin, J.L., 2004. Links between phytoplankton and bacterial community dynamics in a coastal marine environment. Mar. Ecol. 49(1), 163-175.<https://doi.org/10.1007/s00248.003.1057.0>.

Sebastian, M., Pitta, P., Gonzalez, J.M., Thingstad, T.F., Gasol, J.M., 2012. Bacterioplankton groups involved in the uptake of phosphate and dissolved organic phosphorus in a mesocosm experiment with P-starved Mediterranean waters. Envir. Microbiol. 14 (9), 2334-2347. <https://doi.org/10.1111/j.1462-2920.2012.02772.x>

Strickland, J.D.H., Parsons, T.R., 1965. A manual of sea water analysis, In: Stevenson, J.C. (Ed.), Fisheries Research Board of Canada, No. 125, second ed. Roger Duhamel FRSC, Ottawa, Canada, pp. 37-86.

Taga, N., Kobori, H., 1978. Phosphatase activity in eutrophic Tokyo Bay. Mar. Biol. 49 (3), 223-229.<https://doi.org/10.1007/BF00391134>

UNESCO, 1994. Protocols for the Joint Global Ocean Flux Study (JGOFS) Core Measurements. Manual and Guides No 29. pp. 97-100.

Vidal, M., Duarte, C.M., Agusti, S., Gasol, J.M., Vaquer, D., 2003. Alkaline phosphatase activities in the central Atlantic Ocean indicate large areas with phosphorus deficiency. Mar. Ecol. Prog. Ser. 262, 43-53.<https://doi.org/10.3354/meps.262043>.

Wurl, O., 2009. Practical guidelines for the analysis of sea water. CRC Press, Boca Raton. pp.143-178.

Yucel, N., 2018. Spatio-temporal variability of the size fractionated primary production and chlorophyll in the Levantine Basin (northeastern Mediterranean). Oceanol. <https://doi.org/10.1016/j.oceano.2017.12.003>.

## Figure legends

**Figure 1** Spatio-temporal variations in a. chlorophyll and b. phaeophytin off Trivandrum

**Figure 2** Spatio-temporal variations in phosphatase producing bacteria (PPB-MPN) off Trivandrum

**Figure 3** Spatio-temporal variations in alkaline phosphatase activity (APA) off Trivandrum




Fig. 1a




Fig.1b




Fig. 2




Fig. 3

**Table 1 Range and average of physico-chemical and biological parameters in whole water column (5 days, 4 time interval, 4 depths; n=80)**

<b>Parameter</b>	<b>Range</b>	<b>Mean</b>
<b>Temperature (°C)</b>	27.2-28.7	28±0.57
<b>Salinity</b>	33.07-35.46	34.9±0.57
<b>pH</b>	8.19-8.35	8.31±0.04
<b>DO (ml L<sup>-1</sup>)</b>	1.19-4.68	3.47±0.64
<b>NO<sub>2</sub>-N (µM)</b>	ndl-0.795	0.16±0.21
<b>NO<sub>3</sub>-N (µM)</b>	0.681-20.026	5.32±4.68
<b>PO<sub>4</sub>-P (µM)</b>	0.093-2.37	0.27±0.29
<b>Chl <i>a</i> (µg L<sup>-1</sup>)</b>	0.71-2.90	1.39±0.5
<b>Phae (µg L<sup>-1</sup>)</b>	0.15-1.29	0.5±0.3
<b>TBC (cells L<sup>-1</sup>)</b>	2.48x10 <sup>8</sup> -1.54x10 <sup>10</sup>	5x10 <sup>9</sup> ±2.85x10 <sup>9</sup>
<b>TVCa (cells L<sup>-1</sup>)</b>	6.37x10 <sup>8</sup> -8.16x10 <sup>9</sup>	3.3x10 <sup>9</sup> ±2x10 <sup>9</sup>
<b>PPB-MPN (cells L<sup>-1</sup>)</b>	ndl-1.5x10 <sup>6</sup>	0.27x10 <sup>6</sup> ±0.46x10 <sup>6</sup>
<b>APA (µM P hr<sup>-1</sup>)</b>	ndl-13.79	2.34±2.71

DO-dissolved oxygen; Chl*a*-Chlorophyll *a*; Phaeo-Pheaophytin TBC- Total bacterial count; TVCa Total viable bacteria-aerobic; PPB-Phosphatase producing bacteria; APA-alkaline phosphatase activity; ndl-non detectable level

**Table 2 Relationships of alkaline phosphatase activity (APA) with environmental parameters**

	<b>Whole water (n=16)</b>	<b>6 hrs (n=20)</b>	<b>12hrs (n=20)</b>	<b>18hrs (n=20)</b>	<b>24hrs (n=20)</b>
<b>Depth</b>	0.616 <sup>b</sup>	-	0.631 <sup>b</sup>	-	0.468 <sup>c</sup>
<b>DO</b>	-0.547 <sup>c</sup>	-	-0.608 <sup>b</sup>	-	-
<b>NO<sub>3</sub>-N</b>	0.637 <sup>b</sup>	-	-	0.625 <sup>b</sup>	-
<b>Chl a</b>	0.670 <sup>b</sup>	-	0.619 <sup>b</sup>	-	-
<b>Phaeo</b>	0.740 <sup>c</sup>	-	0.638 <sup>b</sup>	-	0.491 <sup>c</sup>
<b>TBC</b>	0.561 <sup>a</sup>	-	-	-	-
<b>TVC</b>	0.491 <sup>a</sup>	-	-	-	-
<b>PPB-MPN</b>	-	0.465 <sup>c</sup>	-	-	-

Superscripts denote different significant levels; a-p<0.05; b-p<0.01 and c-p<0.001. DO-Dissolved Oxygen; Temp.-Temperature; Chl a –Chlorophyll; Phaeo-Pheophytin; TBC-Total Bacterial Count; TVC-Total viable count; PPB-Phosphatase producing bacteria; APA-Alkaline phosphatase activity