

Environmental influences on the species diversity, biomass and population density of soft bottom macrofauna in the estuarine system of Goa, west coast of India

Sadanand N Harkantra* & Nimi R Rodrigues

Biological Oceanography Division, National Institute of Oceanography, Dona Paula, Goa 403004, India

(*E-mail: sadanand@darya.nio.org)

Received 12 June 2003, revised 17 February 2004

A total of 58 species were recorded belonging to polychaetes, molluscs, crustaceans and other minor groups in order of species abundance. eighteen species are new to the local fauna that were not reported earlier. The maximum mean species diversity index (Shannon-Wiener), total biomass (wet) and total population density recorded were 2.3 (Z1), 6.7 g/m² (M1) and 703 no./m² (M2) respectively. Significantly higher species diversity was observed at high salinity, fine sand and high sedimentary biochemical parameters of total organic carbon (TOC), total organic nitrogen (TON) and carbon of biopolymeric fraction (C-BPF) sites. Medium grain size sites supported significant high biomass whereas population density showed no significant difference among the sites. The best multiple linear regression model revealed that all the 13 parameters studied were significant influencing parameters on species diversity, biomass and population density with exception of temperature. Among these salinity and TON were the main significant parameters. The combination of significant influencing environmental parameters, % variation and Mallows' *C_p* values varied from sites to biotic parameters. This explained 32-72% of the total variance. The regression model derived from this data helps in detection of these biotic parameters and detection of pollution-induced effects.

[Keywords: Soft bottom macrofauna, species diversity, biomass, population density, linear regression model, Mandovi estuary, Zuari estuary, Goa]

[IPC Code: Int.Cl.⁷ A01]

Introduction

One of the aims of benthic ecologist is to understand the ecological process, which is achieved by examining the interrelationship between environmental parameters and benthic community structure¹⁻⁴, anthropogenic impacts⁵, and modelling of the ecosystem⁶. This is usually accomplished by inferential analysis of empirical data by multivariate techniques and by conducting manipulative experiments¹⁻⁴. Estuaries form an ideal ecosystem to examine such interactions between the abiotic and biotic factors due to their wide range of these parameters¹⁻⁴. Several studies have examined relative influence of environmental parameters on species diversity, biomass and population density of soft bottom macrofauna in temperate estuaries¹⁻⁶ more than the tropical estuarine systems⁷⁻¹⁰. However, magnitude of variation will change at a particular geographical area⁴. There have been some studies on the Goa estuaries soft bottom macrobenthos^{9,10}. These studies were restricted to a few abiotic and biotic parameters and seldom multivariate techniques have been used. The estuaries of Goa are used for fishing, aquaculture, ore transport, harbour development, waste disposal, adjacent land for human

settlement and water recreation^{8,10,11}. Ore refuge run off during monsoon and its impact on marine biota is one of pollution issues in these estuaries¹⁰. Spatio-temporal variations in abiotic and biotic parameters in this estuarine system are affected by tropical southwest monsoon, riverine and tidal flows contributing to the ecological complexity of the ecosystem^{8,10,11}. Hence, a study was conducted (1997-98) at four sites in Mandovi-Zuari twin tropical estuarine system of Goa, central west coast of India, to test the hypothesis that environmental parameters (abiotic parameters) significantly affect the soft bottom macro-fauna such as species diversity (Shannon-Wiener), total wet biomass and total population density (biotic parameters). Best linear multiple regression models and ANOVA tests were used for one year (1997-98) spatio-temporal data at 4 sites in these estuaries. The regression model derived from such studies helps in detection of these biotic parameters and detection of pollution-induced effects by knowing the environmental parameters.

Materials and Methods

The Mandovi and Zuari estuaries of Goa interconnected by Cumbarjua canal and fringed with

mangroves are located on central west coast of India (Fig. 1). The site M1 is at the proximity of Arabian Sea in Mandovi estuary (depth 3.5 m). The salinity varies between 9.1-32.5 psu. The substratum is sandy and the sediment is composed of 81.2% sand, 7.6% silt and 11.1% clay. Site M2 is situated 3 km further upstream from M1 (depth 3 m) and the salinity varies between 1.1-33 psu. The substratum is sandy and sediment is composed of 94.2% sand, 3.5% silt and 2.1% clay. Site Z1 lies upstream from the mouth of Zuari estuary (depth 3.5 m) and the salinity varies between 20.5-33 psu. The substratum is silty sand and sediment is composed of 62.3% sand, 19.2% silt and 18.4% clay. Site Z2 is about 3 kms upstream from Z1 (depth 3.5 m). The salinity varies between 3-33 psu. The substratum is sand-silt-clay and sediment is composed of 35.6% sand, 27.9% silt and 36.5% clay. The average annual rainfall of Goa is about 3000 mm of which nearly 80% occurs during the southwest monsoon period (June-September), while relatively stable conditions prevail during postmonsoon (October-January) and premonsoon (February-May) period.

Four sites (Fig. 1) were sampled monthly from October 1997 to September 1998. Triplicate samples were obtained at each site with a van Veen grab (0.04 m², 15-20 cm penetration). Soft bottom macrofauna were sieved (0.5mm), preserved, assorted as reported earlier^{12,13}. Macrofauna were identified and food and feeding habits of benthic fauna were ascertained¹⁴. Population density was converted into no./m² and biomass was expressed as wet weight g/m² after removing the hard parts. Sediment texture were analysed according to Buchanan¹⁵. Total organic carbon (TOC) and total nitrogen (TON) of sediment were determined using dried sediment sub-samples (5-10 mg) using NCS 2500 elemental analyser (Italy) after acidification with 1 N HCl. Proteins¹⁶, carbohydrates¹⁷, lipids¹⁸ and carbon of biopolymeric fraction¹⁹ (C-BPF) were also estimated. Bottom water was also collected using a Niskin bottle from the same locations for chlorophyll *a*²⁰ (Chl-*a*) and dissolved oxygen²¹ (DO) estimation. Bottom water salinity and pH were measured using a Guildline Autosol (Model 8400 A, USA) and pH meter Model LI 612 (Elico Pvt Ltd, India). All the estimates were made in triplicate.

Shannon-Wiener species diversity index ($H' - \log_2$ base) was calculated according to Pielou²² for months data of different sites. ANOVA tests were carried out for 13 environmental and 3 biotic parameters to test any significant differences among the sites. Further,




Fig. 1—Map of study sites (solid circle)

one-way Tukey's HSD multiple comparison was used when significance was detected ($P < 0.05$). The best multiple linear regression models²³ were used to assess the relative significant influencing environmental parameters on species diversity, total wet biomass and total population density and to construct the predictive models. The regression explaining the greatest amount of variation (R^2) with all the significant parameter coefficients (r) were presented as the best fit based on adjusted R^2 and minimum Mallows' C_p ²⁴ (low value of C_p indicates more accuracy of the regression model). The total data of four sites were first analysed to give the regression model. However, this did not give any significant regressions and variation was very low. Moreover, ANOVA tests revealed significant differences in environmental parameters among the sites. Hence, it was decided to perform these analyses on the data set of each site. All the raw data were normalized by log₁₀ transformation before any statistical analyses. Minitab-Version 8.3 was used for these analyses²⁵.

Results

Environmental parameters

Table 1 shows annual mean of 13 environmental parameters studied at all the sites. All the environmental parameters varied significantly among the sites, except pH and Chl-*a* (ANOVA, $P < 0.0001$, Tukey's test Table 1). Temperature measurements ranged from 27.2-30.4 °C, DO 3.9-4.4 ml/l, pH 7.3-8.2 and Chl-*a* 0.9-1.7 µg/g. Salinity varied between 12.7 (M2) to 28.8 (Z1) and was significantly higher at sites Z1 in Zuari estuary (ANOVA

$P < 0.0001$, Tukey's test, Table 1). Mean grain size varied from 0.03 (Z2) to 0.3 (M2) and was significantly finer at sites in Zuari than in Mandovi estuary (ANOVA, $P < 0.0001$, Tukey's test Table 1). Sediment sorting varied from 0.3 (Z2) to 0.7 mm (M2) and showed well sorted to poorly sorted sediments. The muddy regions in Zuari estuary varied from silty sand to sand-silt-clay whereas sandy region in Mandovi varied from medium to fine sand. Sedimentary biochemical parameters such as TOC, TON, proteins, carbohydrates, lipids and C-BPF fraction showed wide range of variations (Table 1). TOC varied from 2587 (M1) to 17198 (Z2), TON 334.1 (M2) to 2015.5 (Z2), protein 8.4 (M1) to 18.3 (Z2), carbohydrate 309.4 (M1) to 867.1 (Z2), lipids 75 (Z2) to 140 (M1) and C-BPF 232.4 (M1) to 422.4 (Z1) $\mu\text{g/g}$. Relatively, these values were significantly higher at sites in Zuari than in Mandovi estuary (ANOVA, $P < 0.0001$, Tukey's test, Table 1).

Benthic community structure

A total of 144 grab samples (0.04 m^2) were sorted equivalent to an area of 5.8 m^2 . This yielded 2344 individuals belonging to polychaetes, molluscs, crustaceans and other minor groups in order of species abundance. Polychaetes formed the dominant taxa followed by molluscs and crustaceans. A total of

58 species were recorded of which 18 species were new to the local fauna that were not reported earlier. These included 24 species of polychaetes, 13 species of bivalves, 1 species of gastropod, 12 species of crustaceans, 1 species of fish and 7 other species that included Sipuncula, Nemertea, Anthozoa, Echiura, Nematoda and Coelenterata. The benthic fauna largely composed of the deposit feeding polychaetes *Prionospio pinnata* Ehlers and *Clymene annandalei* Southern at sites in Zuari estuary and the filter feeding bivalve *Meretrix casta* (Chemnitz) and the carnivorous polychaete *Nereis capensis* Wiley at sites in Mandovi estuary. The pattern of species succession and site wise species density has been described in detail elsewhere²⁶.

The annual mean (with standard deviation) of benthic species diversity index (Shannon-Wiener), total wet biomass and total population density at different sites is given in Table 1. The species diversity values varied between 1.5 (M2)-2.2 (M1), total wet biomass 1.7 (M2)-6.7 g/m^2 (M1) and total population density 610 (Z1)-687nos/ m^2 (M2). Significant high species diversity was observed at site M1 due to high salinity and fine sand content and at site Z1 due to high sedimentary TOC, TON, C-BPF. Species diversity and biomass varied significantly among the sites However, population density showed

Table 1—Annual mean (\bar{X}) and standard deviation ($\pm\text{SD}$) of 13 environmental and 3 biotic parameters at different sites. Asterisk indicates variables significant level for ANOVA tests among the sites. The superscripts refer to Tukey's HSD test, where values with same letter in each row are not different at the 0.05 significant level. **** $P < 0.0001$, Ns- Not significant. N=36.

Variables	Sites			
	M1	M2	Z1	Z2
	$\bar{X} \pm\text{SD}$	$\bar{X} \pm\text{SD}$	$\bar{X} \pm\text{SD}$	$\bar{X} \pm\text{SD}$
Temp.****	27.2 \pm 1.0 ^a	29.0 \pm 1.1 ^b	29.8 \pm 1.1 ^b	30.4 \pm 1.1 ^b
Sal.****	21.9 \pm 1.6 ^b	12.7 \pm 3.1 ^a	28.8 \pm 1.1 ^c	20.1 \pm 1.9 ^b
DO ₂ ****	4.3 \pm 1.1 ^b	5.0 \pm 1.5 ^c	3.9 \pm 1.1 ^a	4.4 \pm 1.1 ^b
pH(Ns)	7.0 \pm 1.1	6.8 \pm 1.1	7.3 \pm 1.1	7.0 \pm 1.1
Chl- <i>a</i> (Ns)	1.0 \pm 2.8	0.9 \pm 7.0	1.7 \pm 1.8	1.6 \pm 2.5
M.gr.****	0.2 \pm 0.6 ^b	0.3 \pm 0.8 ^a	0.1 \pm 0.7 ^c	0.03 \pm 0.9 ^d
Sd.s.****	0.5 \pm 0.9 ^b	0.7 \pm 0.9 ^a	0.5 \pm 0.8 ^b	0.3 \pm 0.9 ^c
TOC****	2587.0 \pm 3.5 ^a	2759.3 \pm 2.2 ^a	11697.7 \pm 2.5 ^b	17198.0 \pm 1.7 ^c
TON****	337.7 \pm 2.1 ^a	334.1 \pm 1.8 ^a	1174.0 \pm 1.8 ^b	2015.5 \pm 2.2 ^c
Pr.****	8.4 \pm 2.3 ^a	8.5 \pm 1.5 ^a	15.3 \pm 1.5 ^c	18.3 \pm 1.3 ^c
Carb.****	309.4 \pm 2.9 ^a	355.9 \pm 1.8 ^a	615.0 \pm 2.0 ^b	867.1 \pm 1.5 ^c
Lip.****	140.0 \pm 1.1 ^b	117.2 \pm 1.4 ^b	88.4 \pm 1.9 ^a	75.0 \pm 2.1 ^a
C-BPF****	232.4 \pm 1.6 ^a	236.1 \pm 1.4 ^a	305.2 \pm 1.7 ^b	422.4 \pm 1.3 ^c
H'****	2.2 \pm 0.6 ^c	1.5 \pm 0.7 ^a	2.3 \pm 0.7 ^c	1.9 \pm 0.4 ^b
B****	6.7 \pm 0.4 ^b	1.74 \pm 0.2 ^a	1.8 \pm 0.1 ^a	2.7 \pm 0.1 ^a
P(Ns)	687.1 \pm 1.6	703.3 \pm 3.5	610.1 \pm 0.9	615.0 \pm 0.1

Temp. -Temperature ($^{\circ}\text{C}$), Sal.-Salinity (psu), DO-Dissolved oxygen (ml/l), Chl-*a*- Chlorophyll-*a* ($\mu\text{g/g}$), M.gr.- Mean grainsize (mm), Sd.s.- sediment sorting (mm), TOC-Total organic carbon ($\mu\text{g/g}$), TON -Total organic nitrogen ($\mu\text{g/g}$), Pr.- Protein ($\mu\text{g/g}$), Carb.- Carbohydrates ($\mu\text{g/g}$), Lip.- Lipids ($\mu\text{g/g}$), C-BPF-Carbon of biopolymeric fraction ($\mu\text{g/g}$).

no significant differences (ANOVA, $P < 0.0001$, Tukey's test, Table 1). Relatively, significant high biomass was also observed at site M1 which had fine grain size.

The best multiple linear regression model

All the 13 environmental variables significantly ($P < 0.05-0.0001$) influenced the species diversity index, biomass and population density at all the sites for the significant ($P < .001-0.0001$) best multiple linear regression model except for temperature (Table 2). This explained 32-72% of the total variation (Table 2). No significant best regression fit could be obtained at site Z2 for species diversity. Water properties like salinity and dissolved oxygen were the significant ($P < 0.05-0.0001$) influencing parameters in terms of ability to explain biotic parameters at all the sites (Table 2). Similarly, sediment properties like mean grain size, sediment sorting and sedimentary biochemical variables were the significant ($P < 0.05-0.0001$) influencing environmental parameters to biotic parameters at all the sites except for species diversity at site Z2 (Table 2). Salinity and TON were the significant dominant abiotic parameters in influencing the biotic parameters in view of their higher frequency of occurrence in combination of two or three variables (Table 2). Although we observed significant regression fit ($P < 0.001$, $P < 0.0001$, Table 2), their

combination of significant influencing parameters, % variation (R^2) and Mallows' C_p values varied from sites to biotic parameters.

The % variation (R^2) of best multiple regression model (Table 2) varied between 37-39% for species diversity, 41-70% for biomass and 32-72% for population density at all the sites. Similarly C_p values varied between -3.2-4.0, 2.0-29.8 and 5.1-71.9 for species diversity, biomass and population density respectively (Table 2). The predictive best multiple linear model derived from abiotic parameters also changed from sites to biotic parameters (Table 3)

Discussion

We have examined 13 environmental parameters and their relative influence on the species diversity, biomass and population density of soft bottom fauna. Salinity was a significant parameter to explaining the variance in especially species diversity at site M1 ($P < 0.001$, Table 2) and biomass at all the sites ($P < 0.0001$, Table 2). Wide fluctuation in salinity was mainly brought about by southwest monsoon, land runoff, riverine and tidal flow¹¹. These large variations in salinity in an estuary increases physiological stress, which can result in the reduction in the number of species and restrict the species distribution depending on the species salinity tolerance thresholds¹⁻⁴. For example, some euryhaline species like *Thalassema* sp. (echiuran worm) was

Table 2—Best multiple linear regression (BMR) model for species diversity (H'), total wet biomass (B), and total population density (P) of 13 environmental variables at different sites. Asterisk indicates variables and BMR significant level. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$, Ns-no significance. N=36.

variables	Sites											
	M1			M2			Z1			Z2		
	H'	B	P	H'	B	P	H'	B	P	H'	B	P
Temp.												
Sal.	***	****			****			****			***	
DO	***						***		****			**
pH							**					
Chl- <i>a</i>												**
M.gr.			***	****								
Sd.S.								***	****		****	
TOC		****		*		***						
TON		****	***		**	****			****			***
Pr.	*								****			
Carb.						***	***					
Lip.				***								
C-BPF.			****		****						****	
BMR	****	****	****	****	****	****	****	****	****	Ns	****	***
Ad.R ²	0.39	0.7	0.59	0.37	0.41	0.41	0.39	0.61	0.72		0.51	0.32
C_p	-3.2	3.2	13.9	-3.2	2	11.8	0.4	5.7	5.1		29.8	71.9

Temp. -Temperature (°C), Sal. -Salinity (psu), DO -Dissolved oxygen (ml/l), Chl-*a*-Chlorophyll-*a* (µg/g), M.gr.- Mean grainsize mm, Sd.s.- Sediment sorting (mm), TOC-Total organic carbon (µg/g), TON -Total organic nitrogen (µg/g), Pr.- Protein (µg/g), Carb.- Carbohydrates (µg/g), Lip.- Lipids (µg/g), C-BPF.- Carbon of biopolymeric fraction (µg/g).

Table 3—Best significant linear multiple regression model for predicting the community structure from the environmental parameters at different sites. Ns-No significant regression, N=36.

M1	H' = 0.27-4.46 DO+ 5.14 Sal. + 0.49 Pr. B = 0.28 +2.06 Sal. + 2.65 TOC-4.14 TON P = -1.94 + 0.78 M.gr.-1.80 TON + 4.11 C-BPF
M2	H = -8.27 + 1.30 M.gr. + 0.57 TOC + 2.47 Lip. B = 3.54-0.40 Sal. + 0.34 TON - 1.18 C-BPF P = 1.44 + 3.58 TOC-5.98 TON + 2.54 Carb.
Z1	H' = 13.5 -5.69 DO - 5.93 pH - 0.95 Carb. B = -2.95 + 2.95 Sal. - 0.17 Sd.s. P = 9.98 - 12.3 DO + 1.31 Sd.s. + 2.64 TON - 6.07 Pr.
Z2	H' = Ns B = 1.06 + 0.41 Sal. - 0.69 TOC + 0.99 C-BPF P = 13.0 - 0.75 Chl- a - 7.14 pH - 0.35 M.gr.

Temp. -Temperature (°C), Sal. -Salinity (psu), DO -Dissolved oxygen (ml/l), Chl-*a*-Chlorophyll-*a* (µg/g), M.gr.-Mean grain size (mm), Sd.s.-Sediment sorting (mm), TOC-Total organic carbon (µg/g), TON -Total organic nitrogen (µg/g), Pr.- Protein (µg/g), Carb.- Carbohydrates (µg/g), Lip.- Lipids (µg/g), C-BPF.- Carbon of biopolymeric fraction (µg/g).

only found in high salinity site (M1), whereas oligohaline species like *M. casta* (bivalve) was only observed in low salinity sites (M2) in Mandovi estuary. Such a distribution and the species succession has been described elsewhere²⁶. This indicates that salinity acts as a physiological barrier for oligo- and euryhaline species due to the functional physiology of the species¹⁻⁴. This parameter is more critical in the early larval stages of benthos¹⁻⁴. Hence, species diversity reduced in low salinity sites and increased at high salinity sites (Table 1) supporting earlier observations¹⁻⁴. We also observed relatively significant ($P < 0.0001$) high biomass in high salinity site (M1) (Table 1).

Dissolved oxygen variations which are dependant on the mixing process 11 was a significant parameter in explaining especially species diversity (M1 and Z1, $P < 0.001$) and population density (Z2, $P < 0.0001$, Table 2) and suggests oxygen threshold²⁷. Typically, macrofauna exhibit reduced diversity and extreme dominance in oxygen minimum zone²⁷. However, we did not observe such phenomenon. Oxygen was not a significant influencing parameter with respect to biomass (Table 2).

Some portion of sedimentary biochemical parameters (C-BPF) which forms the food for benthic organisms²⁸ was seen as a significant ($P < 0.05$ - 0.0001) parameter explaining the variation in species diversity (M1, M2 and Z1) biomass (M1, M2 and Z2) and population density at all the sites (Table 2). This is mainly attributed to the food and feeding

relationship²⁸. Bottom water Chl-*a* which forms food for the benthic organisms²⁹ largely depends on overlying water column productivity²⁹. This parameter was significant ($P < 0.05$) in explaining the population density only at site Z1 (Table 2). The same was true for pH at site Z1, with respect to species diversity, which is dependent on water quality¹¹.

Mean grain size and sediment sorting distribution which is a function of hydrodynamic regime³⁰ played a significant ($P < 0.001$, $P < 0.0001$) role in explaining species diversity, biomass and population density at all the sites (Table 2). Relatively finer sediments supported significant ($P < 0.0001$) high species diversity at sites M1 and Z1 (Table 1) corroborating earlier findings and indicating wider niches³¹. Well sorted sediments usually contain a small range of grain sizes and interstitial spaces. Such areas may provide fewer niches than poorly sorted sediment areas and which contain a less diverse fauna as noticed at site M2 (Table 1), thus supporting the earlier findings³¹. Absence of significant regression fit at some site for biotic parameters (Table 2) partly reflects the influence of unmeasured environmental parameters, such as hydrodynamic and sediment transport process, that affect the distribution of both sediments and fauna³⁰. The other possible reason is that it may be linked with other unmeasured environmental parameters and biotic factors in the absence of a significant relationship.

Multivariate statistical methods can extract information from data sets containing larger amounts of variance, considering simultaneously the interrelationships of several influential variables³²⁻³³. The major ecological assumption for the development of this model was that the benthic community structure fluctuates greatly in relation to the environmental conditions *in situ* rather than mortality, inter-specific competition or predation^{32,33}. Further, the influence of abiotic factors acting on the benthic community prior to sampling or the duration there of were not considered³². It is clear from an examination of the R^2 values that the predictor variable subsets selected, did not account for all the variance in the dependant variable. Several reasons may be attributed to this^{23,32,33}. Firstly, the important variables with a strong influence were omitted due to multicollinearity and the procedure used to fit the linear regression model assumes that predictive variables are measured without errors³³. Therefore, any error in benthic community structure would certainly introduce variance in the model that cannot be accounted for

any independent variable^{32,33}. The accuracy did not improve by incorporating additional variables. Fitting hypothetical models to such a data-set is a compromise unless the variables and relationships are correctly chosen^{23,32,33}.

All the 13 environmental parameters except temperature were shown to be significant influencing parameters on species diversity, biomass and population density that explained 32-72% of the total variance. Among these, salinity and TON were the main significant factors. Unexplained variations and absence of significant regression fit suggests the influence of unmeasured environmental parameters and biotic factors. Hence, *in situ* manipulative experiments are needed to understand the biotic factors in this area. The model shown here is a pure representation of the available data set and interpretations are limited to the study site and period. The regression model derived from this data (Table 3) helps in detection of these biotic parameters and detection of pollution-induced effects by knowing the environmental parameters.

Acknowledgement

We thank Director, NIO, Goa, for encouragement and facilities. The work was carried out under the Indo-US Project on Trophic Dynamics funded by Office of Naval Research, Washington. We also thank Dr. S.G. Dalal for useful discussion on statistical analysis, Dr. D. Chandramohan and other colleagues for help during various stages of this work. N.R.R. thanks CSIR, for award of a Senior Research Fellowship. This is NIO contribution No. 3869

References

- Edgar G J & Barrett N S, Benthic macrofauna in Tasmanian estuaries, Scale of distribution and relationship with environmental variables, *J Exp Mar Biol Ecol*, 270 (2002) 1-24.
- Brown S S, Gastona G R & Rakocinski C F, Effect of sediment contaminants and environmental gradients on macrobenthic community trophic structure in Gulf of Mexico estuaries, *Estuaries*, 23 (2000) 411- 424.
- Mannino A & Montagna P A, Small-scale spatial variation of macrobenthic community structure, *Estuaries*, 20 (1997) 159-173.
- Zajac R N & Whitlatch R B, Response of estuarine infauna to disturbance II Spatial and temporal variation of succession, *Mar Ecol Prog Ser*, 10 (1982) 25-27.
- Frouin P, Effects of anthropogenic disturbance of tropical soft bottom benthic communities, *Mar Ecol Prog Ser*, 194 (2000) 39-53.
- Longhurst A R, Ecological models in estuarine management, *Oceanogr Mangmt*, 4 (1978) 287-302.
- Harkantra S N & Parulekar A H, Community structure of sand dwelling macrofauna of an estuarine beach in Goa India, *Mar Ecol Prog Ser*, 30 (1985) 291-294.
- Parulekar A H, Dhargalkar V K & Singbal S Y S, Benthic studies in Goa estuaries: Part III-Annual cycle of macrofaunal distribution, production and trophic relations, *Indian J Mar Sci*, 9 (1980)189-200.
- Harkantra S N, Benthos of the Kali estuary, Karwar, *Mahasagar-Bull Natn Isnt Oceanogr*, 8 (1975) 53-58.
- Parulekar A H, Ansari Z A & Ingole B S, Effect of mining activities on clam fisheries and bottom fauna of Goa estuaries, *Proc Indian Acad Sci (An Sc)*, 95 (1986) 325-339.
- Qasim S Z & Sen Gupta R, Environmental characteristics of the Mandovi- Zuari estuarine system in Goa, *Estuar Coast Shelf Sci*, 13 (1981) 557- 578.
- Eleftheriou A & Holme N A, Macrofauna techniques, in : *Methods for the study of marine benthos* edited by Holme N A & McIntyre A D, (Blackwell Scientific Publications, Oxford) 1984, pp.140-216.
- Harkantra S N & Parulekar A H, Soft sediment dwelling macroinvertebrates of Rajapur Bay, central west coast of India, *Indian J Mar Sci*, 23 (1994) 31-34.
- Fauchald K & Jumars P A, The diet of worms: A study of polychaete feeding guild, in: *Oceanography and marine biology*, Vol.16, edited by M Barnes (Allen and Unwin, London) 1979, 193-284.
- Buchanan J B, Sediment analysis, in : *Methods for the study of marine Benthos*, edited by Holme N A & McIntyre A D, (Blackwell Scientific Publications, Oxford) 1984, pp. 41-65.
- Lowry O H, Rosenbrough N J, Farr A L & Randall K J, Protein measurements with folin phenol reagent, *J Biol Chem*, 193 (1951) 265- 275.
- Kochert G, Carbohydrate determination by the phenol-sulphuric acid method, in: *Handbook of phycological method: Physiological and biochemical methods*, edited by Hellebesr J A & Craigie J S (Cambridge University Press, Cambridge) 1978, pp. 95-97.
- Parsons T R, Maita Y & Lalli C H, *A manual of chemical and biological methods for sea water analyses*, (Pergaman Press, Oxford) 1984, pp. 75-80.
- Fabiano M, Danovaro R & Fraschetti S, A three year time series of elemental and biochemical composition of organic matter in sub-tidal sandy sediments of the Ligurian Sea (Northwestern Mediterranean), *Cont Shelf Res*, 15 (1995) 1453-1469.
- Lorenzen C J, Determination of chlorophyll and phaeopigment: Spectrophotometric equation, *Limnol Oceanogr*, 12 (1966) 343- 346.
- Strickland J D & Parsons T R, *A practical handbook of seawater analysis*, *Bull Fish Res Board Canada*, Ottawa, No.167, 1972, pp.310.
- Pielou E C, *Ecological diversity*, (Wiley Interscience Publications, London) 1975, pp.165.
- Draper N R & Smith H, *Applied regression analysis*, 2 nd edition, (Wiley and Sons, New York) 1981, pp. 709.
- Helsel D R & Hirsch R M, *Statistical methods in water resources*, (Elsevier Science Publishers, Amsterdam) 1992, pp. 522.
- Minitab, Inc. *Minitab reference manual: release 8.2 for PC* (State College, Pennsylvania, USA) 1991, pp. 234.
- Harkantra S N & Rodrigues N, Pattern of species succession of soft bottom macrofauna in the estuaries of Goa, west coast of India, *Curr Sci*, 85 (2003) 1458-1464.

- 27 Diaz R J, & Rosenberg R, Marine benthic hypoxia: a review of its ecological effects and the behavioral response benthic macrofauna, *Oceanogr Mar Biol*, 33 (1995) 245-303.
- 28 Marsh A G & Tenore K R, The role of nutrition in regulating the population dynamics of opportunistic surface deposit feeder in a mesohaline community, *Limnol Oceanogr*, 35 (1990) 710 -734.
- 29 Christensen H & Kannevorff R , Sedimenting phytoplankton as major food sources for suspension and deposit feeders in the fresund, *Ophelia*, 24 (1985) 223-244.
- 30 Snelgrove P V R & Butman C A, Animal-sediment relationships revisited: Causes versus effect, *Oceanogr Mar Biol*, 32 (1994)111-177.
- 31 Coleman N, Gason A S H & Poore G C B, High species richness in shallow marine waters of south-east Australia, *Mar Ecol Prog Ser*, 154 (1997) 17-26.
- 32 Clark B M, Bennett B A & Lamberth S J, Temporal variations in surf zone fish assemblages from False Bay South Africa, *Mar Ecol Prog Ser*, 131(1996), 35-47.
- 33 Wiesberg S, *Applied linear regression*, (John Wiley and Sons, New York) 1985, pp.323.