

**Export fluxes of dissolved inorganic carbon to the Northern Indian Ocean from the Indian monsoonal rivers**

**Moturi S. Krishna<sup>1</sup>, Rongali Viswanadham<sup>1</sup>, Mamidala H. K. Prasad<sup>1</sup>, Vuravakonda R. Kumari<sup>1</sup>, Vedula V. S. S. Sarma<sup>1</sup>**

<sup>1</sup>CSIR-National Institute of Oceanography, Regional Centre, Visakhapatnam, 530017, India  
*Correspondence to:* M. S. Krishna (moturi@nio.org)

**Abstract**

Rivers are an important source of dissolved inorganic carbon (DIC) to the adjacent coastal waters. In order to examine the spatial variability in the distribution and major sources of DIC in the Indian monsoonal rivers and to quantify their export flux to the north Indian Ocean, 27 major and medium rivers were sampled during the discharge period. Significant spatial variability in concentrations of DIC (3.4 – 73.6 mg l<sup>-1</sup>) was observed and it is attributed to spatial variations in the precipitation pattern, size of rivers, pollution, and lithology of the catchments. The stable isotopic composition of DIC ( $\delta^{13}\text{C}_{\text{DIC}}$ ) also showed strong spatial variability (-13.0 to -1.4‰) in the Indian monsoonal rivers with relatively depleted  $\delta^{13}\text{C}_{\text{DIC}}$  values in rivers of the northwestern of India (-11.1±2.3‰) and enriched values in the southeastern India (-3.5±2.3‰). Results of the least square linear regression models of Keeling and Miller-Tans plots indicated that chemical weathering of carbonate and silicate minerals by soil CO<sub>2</sub> is the major source of DIC in the Indian monsoonal rivers. Spatial variability in the deviation of  $\delta^{13}\text{C}_{\text{DIC}}$  from the approximated  $\delta^{13}\text{C}$  of source may probably due to dominant autotrophic production in rivers of the southeast region whereas, heterotrophic decomposition of organic matter largely influences in the other Indian monsoonal rivers. It is estimated that the Indian monsoonal rivers annually export ~10.3 Tg of DIC to the northern Indian Ocean, of which the major fraction (75%) enters into the Bay of Bengal and the remaining reaches to the Arabian Sea. This is consistent with the freshwater flux which is three times higher to the Bay of Bengal (~378 km<sup>3</sup> yr<sup>-1</sup>) than to the Arabian Sea (122 km<sup>3</sup> yr<sup>-1</sup>). Despite discharge from the Indian monsoonal rivers account for only 1.3% of the global freshwater discharge, they disproportionately export 2.5% of the total DIC export by the world major rivers. Despite rivers from the SW region of India export an order of magnitude lower DIC (0.3 Tg yr<sup>-1</sup>) than the rivers from other regions of India, the highest yield of DIC was found in the former and it is attributed to intense precipitation (~3000 mm), favorable natural vegetation of tropical moist deciduous and tropical wet evergreen and semi-evergreen forests, tropical wet climate, high soil organic carbon and the dominance of red loamy soils in catchments of the rivers from SW region.

*Keywords:* dissolved inorganic carbon, export flux, Indian rivers, Bay of Bengal, Arabian Sea, North Indian Ocean

## 1. Introduction

Dissolved inorganic carbon (DIC) is one of the major constituent of carbon species in rivers. DIC in rivers mainly originates from the geogenic (weathering of carbonate and silicate rocks) and biogenic (decomposition of organic matter in soils) sources (Meybeck, 1987; Mook and Tan, 1991; Gaillardet et al., 1999, Dessert et al., 2001; Viers et al., 2007; Raymond et al., 2008; Tamooh et al., 2013). The former consumes atmospheric carbon dioxide (CO<sub>2</sub>) while the latter releases CO<sub>2</sub> fixed by the terrestrial plants. In addition to these major sources in the catchment, DIC is also contributed by various physical and biological processes within the rivers. For instance, heterotrophic decomposition of organic matter, photo-oxidation of dissolved organic carbon (DOC), autotrophic respiration and dissolution of atmospheric CO<sub>2</sub> contribute DIC to rivers. On the other hand, autotrophic production by aquatic plants (photosynthesis) and evasion of CO<sub>2</sub> to atmosphere with draw DIC from rivers. All these processes in the catchments and within the rivers are strongly coupled to atmospheric CO<sub>2</sub> because they act as either sink or source of atmospheric CO<sub>2</sub> (e.g. Berner et al., 1983; Mook and Tan, 1991; Gaillardet et al., 1999; Richey et al., 2002). The DIC in rivers and its export to the coastal oceans is thus intimately linked to the global carbon cycle (Campeau et al., 2017)

Riverine export fluxes of DIC to coastal regions of the world oceans have been estimated on the global (Gaillardet et al., 1999) and regional scales (Richey et al., 2002; Wallin et al., 2013; Crawford et al., 2014; Campeau et al., 2014; Kokic et al., 2015) to understand the component of DIC in the global carbon budget. Annual export flux of DIC from the world major river systems to the global ocean has been estimated as ~327 - 385 Tg (1Tg=10<sup>12</sup>g) (Ludwig et al., 1998; Meybeck and Vorosmarty, 1999). However, many of the regional studies on DIC export fluxes were limited only to the major river systems (e.g. Gaillardet et al., 1999; Joesoef et al., 2017), for example, the Mississippi (Raymond and Cole, 2003; Raymond et al., 2008; Cai et al., 2008), Changjiang and Pearl (Cai et al., 2008) and Congo (Wang et al., 2013) rivers etc. Regional studies on the riverine export fluxes of DIC are very important for the global carbon cycle and budget as the export fluxes are largely dependent on the hydrological, lithological and environmental conditions, which are highly variable on the regional scales. However, DIC measurements are still lacking in several medium rivers from different regions of the world in general and Asia in particular.

Studies on the sources and export fluxes of DIC from the Indian rivers are very limited. Though DIC measurements were conducted in some Indian estuaries, for example,

Mandovi and Zuari (Sarma et al., 2001), Godavari (Sarma et al., 2011), Cochin (Gupta et al., 2009; Bhavya et al., 2018), Hooghly (Mukhopadhyay et al., 2002; Samanta et al., 2015), Mahanadi (Pattanaik et al., 2017) and Chilka (Gupta et al., 2008; Muduli et al., 2013), they were confined only to the internal cycling of DIC and exchange of CO<sub>2</sub> at the air-water interface, but not focused on the sources and export fluxes of DIC. The major sources of DIC in the Indian rivers remain unclear, except only a couple of rivers, Krishna (Das et al., 2005; Laskar et al., 2014) and Ganges (Samanta et al., 2015). Further, the quantity of annual DIC export by the Indian rivers to the coastal regions is unknown. Here, we made an attempt to understand the major sources of DIC in the Indian monsoonal rivers (Fig. 1) using  $\delta^{13}\text{C}_{\text{DIC}}$  as a potential tracer, and to estimate the riverine export flux of DIC to the north Indian Ocean from the Indian subcontinent.

The stable isotopic composition of DIC ( $\delta^{13}\text{C}_{\text{DIC}}$ ) is widely used to identify the major sources of DIC in the aquatic systems (e.g. Singh et al., 2005; Tamooch et al., 2013; Samanta et al., 2015; Zou, 2016). The isotopic composition of DIC originated by dissolution of atmospheric CO<sub>2</sub> is about 0‰ (Coplen et al., 2002) whereas it is about -27 to -26‰ if the DIC is derived from oxidation of organic matter produced by C<sub>3</sub> plants (O’Leary, 1988). The  $\delta^{13}\text{C}$  of DIC generated by carbonic acid (formed by soil CO<sub>2</sub> dissolution) weathering of silicates is about -21 to -17‰ (Solomon and Cerling, 1987) while it is in the range of -10 to -9‰ for carbonate rocks because half of the carbon comes from carbonate rocks (0‰, Land, 1980) during weathering. The weathering of silicate and carbonate minerals yield  $\delta^{13}\text{C}_{\text{DIC}}$  in the range of -8 to -7‰ and -4 to -3‰, respectively, if the carbonic acid formed by the dissolution of atmospheric CO<sub>2</sub>. Though the  $\delta^{13}\text{C}$  of DIC derived from different sources is well separable (Deines et al., 1974), the isotopic fractionation by in-stream physical and biological processes alters the  $\delta^{13}\text{C}$  of DIC source (Fig. 2). For example, photosynthesis and degassing enriches (O’Leary, 1988; Finlay, 2004; Parker et al., 2005, 2010; Polsenaere and Abril, 2012; Venkiteswaran et al., 2014) while the heterotrophic decomposition of organic matter and photo-oxidation of dissolved organic carbon depletes the  $\delta^{13}\text{C}$  of DIC (Opsahl and Zepp, 2001; Finlay, 2003; Waldron et al., 2007; Vahatalo and Wetzel, 2008) (Fig. 2). Though, rivers are generally in disequilibrium with atmospheric CO<sub>2</sub> (Raymond et al., 2013) and emit CO<sub>2</sub> to atmosphere due to oversaturation (Oquist et al., 2009; Campeau et al., 2017), the isotopic equilibration between the DIC and CO<sub>2</sub> in the atmosphere significantly influences the  $\delta^{13}\text{C}_{\text{DIC}}$  in rivers (Abongwa and Atekwana, 2014; Deirmendjian and Abril, 2018) due to selective fluxes of <sup>12</sup>CO<sub>2</sub> and <sup>13</sup>CO<sub>2</sub> at the water-air interface. Hence, the influence of

biogeochemical processes within the rivers must be considered while interpreting the  $\delta^{13}\text{C}_{\text{DIC}}$  results for identification of DIC sources. The main objectives of this study are to (i) identify the major sources of DIC in the Indian monsoonal rivers, (ii) estimate the export flux and yield of DIC to the north Indian Ocean and (iii) examine the major processes in the catchments and within the rivers controlling DIC in the Indian monsoonal rivers.

## **2. Study region, sampling and methods**

### **2.1 Study Area**

The Indian peninsula bifurcates the north Indian Ocean into the Bay of Bengal and the Arabian Sea. Although these two basins occupy the same latitudinal belt, their oceanographic processes were reported to be remarkably different due to higher freshwater flux into the Bay of Bengal ( $1.63 \times 10^{12} \text{ m}^3 \text{ yr}^{-1}$ ) than to the Arabian Sea ( $0.3 \times 10^{12} \text{ m}^3 \text{ yr}^{-1}$ ; Subramanian, 1993; Gauns et al., 2005). The large freshwater influx leads to the formation of a strong vertical salinity stratification in the Bay of Bengal (Varkey et al., 1996) that prevents vertical mixing of nutrient rich sub-surface water with the surface (Prasanna Kumar et al., 2004). As a result, the Bay of Bengal is considered to be relatively less productive (Prasannakumar et al., 2002) than the adjacent Arabian Sea, which is one of the highly productive zones in the world (Madhupratap et al., 1996; Smith, 2001; Barber et al., 2001) due to injection of nutrients into the surface through the seasonal upwelling and convective mixing (Shetye et al., 1994; Madhupratap et al., 1996; Muraleedharan and Prasannakumar, 1996).

Discharge from the Indian monsoonal rivers is largely fed by the monsoon induced precipitation over the Indian subcontinent, which receives >80% of its annual rainfall during the southwest (SW) monsoon period (June-September) (Soman and Kumar, 1990). Though some amount of rainfall occurs during the northeast (NE) monsoon (December-March), it does not generate discharge as it will be stored within the dam reservoirs for domestic, industrial and irrigation purposes. Discharge from the Indian monsoonal rivers mainly occurs during the SW monsoon season (Vijith et al., 2009; Sridevi et al., 2015) hence, these rivers are called as monsoonal rivers. Since the major portion of the annual freshwater discharge occurs only during the SW monsoon, the entire estuary is filled with freshwater (Vijith et al., 2009; Sridevi et al., 2015) during this period. As discharge is small during the rest of the year, the discharge during the SW monsoon (wet period) is considered to be equivalent to the annual discharge of the monsoonal rivers. Based on rainfall intensity, forest cover, vegetation

and soil type in the catchment, rivers sampled in the present study were categorized into 4 groups, namely the northwest (NW), southwest (SW), southeast (SE) and northeast (NE) rivers of India (Fig. 1). The SW region of India is characterized by the intense rainfall during SW monsoon (~3000 mm) following the NE (1000-2500 mm), SE (300-500 mm) and NW (200-500 mm) regions of India (Soman and Kumar, 1990). The SW rivers drain red loamy soils while the NW rivers drain black soils. Except the major rivers Godavari and Krishna, all the rivers reaching Bay of Bengal (NE and SE rivers) drain red loamy and alluvial soils in their upper and lower catchments respectively. The Godavari and Krishna rivers drain black soils in their upper catchment whereas red loamy and alluvial soils in their middle and lower catchments respectively (Geological Survey of India; [www.gsi.gov.in](http://www.gsi.gov.in)). Based on discharge, the monsoonal rivers in this study were divided into two types, namely, the major ( $>150 \text{ m}^3 \text{ s}^{-1}$ ) and medium ( $<150 \text{ m}^3 \text{ s}^{-1}$ ) rivers.

## 2.2 Sample collection

Water samples were collected from the freshwater regions of the estuaries to obtain reliable export fluxes of DIC to the coastal ocean. Samples were collected at 2 to 3 locations to minimize the spatial variability within the freshwater zone of the estuary. Further, to minimize the inter-annual variability in DIC concentrations, sampling was conducted in two different years and the mean was used for export flux estimations. Further, samples were collected in mid-stream of the river using a local mechanized boat to avoid the contamination from river banks.

*In-situ* measurements and sample collection were conducted in 27 rivers of the Indian subcontinent (Fig. 1) during the SW monsoon season of the years, 2011 and 2014. Surface water samples at each location were collected for phytoplankton biomass (Chl-*a*), DIC and dissolved oxygen (DO). Samples for DIC were collected in air-tight crimp-top glass bottles and added poison (mercuric chloride) to arrest the biological activity. DO analysis was carried out at a temporary shore laboratory set up for sample processing after the completion of sampling on each day. Water samples were filtered through GF/F (nominal pore size of  $0.7 \mu\text{m}$ ) under moderate vacuum and stored in liquid nitrogen for Chl-*a* analysis.

## 2.3. Methods

Temperature and salinity at the sampling locations were measured using a conductivity-temperature-density (CTD) profiling system (Sea Bird Electronics, SBE 19 plus, United States of America). Concentration of DO was determined by a Winkler's method

(Carritt and Carpenter, 1966) using an auto titrator (Metrohm, Switzerland) with potentiometric end point detection. The analytical precision of the method was  $\pm 0.07\%$  (RSD). Dissolved oxygen saturation is computed following formulations given by Garcia and Gordon (1992). DIC concentrations in water samples were measured at our Institute laboratory using Coulometer (UIC Inc., USA) connected to an automatic sub-sampling system. Based on the repeated analysis of samples and standards, the precision of the method was  $\pm 0.02 \text{ mg l}^{-1}$ . The certified reference materials (CRM) supplied by Dr. A.G. Dickson, Scripps Institute of Oceanography, USA and internal standards were used to test the accuracy of our DIC measurements and it was found to be within  $\pm 0.2$  to  $0.3\%$ . Potentiometric Gran titration method (Metrohm, Switzerland) was used for determination of pH and total alkalinity and followed the standard operating procedures given by Department of Energy (DOE) (1998).

The stable carbon isotopic composition of DIC in the water was measured on Gas Bench coupled with isotope ratio mass spectrometer (EA-IRMS-Delta V, Finnigan, Germany). 50 ml air-tight bottles with rubber septa were filled with 0.5 ml of high purity ortho-phosphoric acid and purged with high purity helium. About 1 ml of water sample is injected to the bottle and incubated at constant temperature of  $50^{\circ}\text{C}$  for 12 hours. The  $\text{CO}_2$  extracted into the head space is injected to the IRMS through gas bench. The results are expressed relative to conventional standards, that is, Pee Dee belemnite (PDB) limestone for carbon (Coplen, 1996) as  $\delta$  values, defined as:

$$\delta R = [(X_{\text{sample}} - X_{\text{standard}}) / X_{\text{standard}}] \times 10^3 \text{ ‰}$$

where R refers to  $^{13}\text{C}$  and X stands for  $^{13}\text{C}/^{12}\text{C}$ . The high-purity tank of  $\text{CO}_2$  was used as working standard for carbon. These gases were calibrated with IAEA standards. Standard deviation on 20 aliquots of the same sample was lower than  $0.05\text{‰}$  for  $\delta^{13}\text{C}$ . Chlorophyll-*a* (Chl-*a*) on the filter was extracted into di-methyl formamide (DMF) and measured the extract fluorometrically using a spectrofluorophotometer (Varian Eclipse, Varian Electronics., UK) following Suzuki and Ishimaru (1990). Annual mean discharge data of the rivers was taken from Meybeck and Ragu (1995, 1996), Central Water Commission, New Delhi (2006, 2012) and Kumar et al. (2005). Catchment area of the rivers was obtained from Water Resources Information System of India (WRIS, [www.india-wris.nrsc.gov.in](http://www.india-wris.nrsc.gov.in)). Soil organic carbon data was taken from Kishwan et al. (2009) and Sreenivas et al. (2016), and the rainfall data was obtained from Soman and Kumar (1990). Dissolved organic carbon (DOC) data for the Indian rivers was taken from Krishna et al. (2015)

Total export flux of DIC from each river was estimated by multiplying the mean concentrations of DIC at near zero salinity (river end member) with the annual discharge. Spatial variability of DIC concentrations within the river was minimized to a large extent by collecting samples from 2 to 3 locations in each river while the inter-annual variability by collecting samples during discharge periods of two years. However, variability in DIC concentrations within the discharge period results in some uncertainties in our estimations of DIC export fluxes. Time series measurements in the Godavari estuary (our unpublished results) revealed that the variability in DIC concentrations within the discharge period is up to 10%. Therefore, the error associated with our DIC flux estimates may be about 10%. DIC flux normalized by catchment area (yield) was calculated by dividing the total DIC export flux of the river by its catchment area.

### **3. Results**

#### **3.1. Hydrographic characteristics**

Surface water temperatures were higher in rivers from the NE and SE regions (mean  $30.9 \pm 1.2^\circ\text{C}$ ) than the rivers from SW and NW regions ( $27.3 \pm 1.5^\circ\text{C}$ ) of India. Dissolved oxygen saturation varied from as low as 63% to as high as 105%, with a mean saturation of  $90 \pm 11\%$ . The rivers from SW region of India recorded more under-saturation of DO ( $82 \pm 7\%$ ) than the rivers located in the NE ( $89 \pm 15\%$ ), NW ( $93 \pm 3\%$ ) and SE ( $96 \pm 11\%$ ) regions of India. Chlorophyll-*a* (Chl-*a*) concentrations varied broadly from 0.8 to  $7.5 \text{ mg m}^{-3}$ , with relatively higher mean concentrations in rivers of the SE region ( $4.7 \pm 2.5 \text{ mg m}^{-3}$ ) followed by the SW ( $2.8 \pm 0.7 \text{ mg m}^{-3}$ ) regions of India. On the other hand, relatively low Chl-*a* was observed in the medium ( $2.6 \pm 1.3 \text{ mg m}^{-3}$ ) than the major estuaries ( $3.2 \pm 2.1 \text{ mg m}^{-3}$ ).

#### **3.2 DIC concentrations and $\delta^{13}\text{C}_{\text{DIC}}$**

DIC concentrations in the Indian monsoonal rivers widely varied from 3.4 (Bharathapuzha) to  $73.6 \text{ mg l}^{-1}$  (Vellar), with a significant spatial variability (Fig. 3a; Table 1). Highest mean DIC concentration was observed in rivers of the SE region ( $37.4 \pm 6.3 \text{ mg l}^{-1}$ ) while the lowest was found in the SW region ( $5.2 \pm 2.1 \text{ mg l}^{-1}$ ) of India. Intermediate values were found in rivers of the NW ( $28.4 \pm 8.9 \text{ mg l}^{-1}$ ) and NE ( $17.1 \pm 6.2 \text{ mg l}^{-1}$ ) regions of India. DIC concentrations were found to be similar in the major ( $22.7 \pm 13.6 \text{ mg l}^{-1}$ ) and medium ( $21.1 \pm 13.2 \text{ mg l}^{-1}$ ) rivers (homoscedastic Student's t-test;  $p=0.76$ ). Mean DIC concentration found in this study ( $21.4 \pm 16.3 \text{ mg l}^{-1}$ ) is similar to those observed earlier in the major river systems of India (Brahmaputra; Singh et al., 2005) and elsewhere in the world, for example,

British rivers (Jarvie et al., 2017) and Swedish rivers (Campeau et al., 2017). However, DIC concentrations in the present study are higher than the global mean DIC (10.3 mg l<sup>-1</sup>, Meybeck and Vorosmarty, 1999) (Table 1), but lower than those reported in the rivers draining into the Gulf of Trieste (N Adriatic; 37-66 mg l<sup>-1</sup>, Tamse et al., 2014).

The  $\delta^{13}\text{C}_{\text{DIC}}$  varied from -13.0 to -1.4‰, with a significant spatial variability (Fig. 3d; Table 1) in the rivers sampled. Relatively depleted  $\delta^{13}\text{C}_{\text{DIC}}$  values were observed in rivers of the NW region (-11.1±2.3‰) while enriched  $\delta^{13}\text{C}_{\text{DIC}}$  was found in rivers of the SE region (-3.5±2.3‰) of India (Fig. 3d). The  $\delta^{13}\text{C}_{\text{DIC}}$  values found in this study are well within the range of values reported earlier in rivers of India (Das et al., 2005) and elsewhere in the world, for example, Swedish streams (-27.6 to -0.6‰; Campeau et al., 2017) and rivers from Italy and Slovenia (-12.8 to -7.7‰, Tamse et al., 2014).

### 3.3. Export fluxes and yield of DIC

Annual export flux of DIC to the coastal ocean from the individual rivers varied broadly from 0.01 Tg (Chalakuudi) to as high as 2.33 Tg (Krishna) (Fig. 3b; Table 1). Among the rivers sampled, rivers of the NE region of India export higher DIC (6.52 Tg yr<sup>-1</sup>) while the lowest was found from rivers of the SW region (0.24 Tg yr<sup>-1</sup>) (Table 1). The Indian monsoonal rivers together export about 10.32 Tg yr<sup>-1</sup> of DIC to the northern Indian Ocean, of which 7.81 Tg (75%) enters into the Bay of Bengal and the remaining into the Arabian Sea (2.51 Tg). The yield of DIC ranged from 2.8 (Bharathapuzha) to 20.7 g m<sup>-2</sup> yr<sup>-1</sup> (Baitarani) (3c; Table 1), excluding the exceptionally high yield of 119 g m<sup>-2</sup> yr<sup>-1</sup> from Haldia river. The mean yield was found to be more or less similar in rivers from all the four regions of India, i.e, NW (8.4 g m<sup>-2</sup> yr<sup>-1</sup>), SW (8.8 g m<sup>-2</sup> yr<sup>-1</sup>), SE (6.6 g m<sup>-2</sup> yr<sup>-1</sup>) and NE (7.7 g m<sup>-2</sup> yr<sup>-1</sup>) regions. Despite the export flux of DIC is lowest from rivers of the SW region (0.24 Tg yr<sup>-1</sup>), interestingly, the yield from rivers of this region is on par with (even slightly higher than) the other Indian monsoonal rivers (Table 1; Fig. 3b&c). Yields of DIC found in this study are similar to those found earlier in rivers elsewhere in the world (Huang et al., 2012).

## 4. Discussion

### 4.1 Distribution of DIC in the Indian monsoonal rivers

Distribution of DIC in the Indian monsoonal rivers showed large spatial variability, with the lowest values in rivers from the SW region of India (Fig. 3a). DIC concentrations in rivers are known to be influenced by the intensity of precipitation over the catchment, basin lithology (Giesler et al., 2013; Lofgren et al., 2014), length of the fluvial network (Hotchkiss



et al., 2015) and in-stream physical and biological processes (Mook and Tan, 1991; Raymond et al., 2008). The spatial distribution of rainfall over the Indian subcontinent ([www.imd.gov.in](http://www.imd.gov.in)) shows that the SW region receives the highest annual rainfall (~3000 mm) than the rest of India (Soman and Kumar, 1990).

The intense precipitation over the SW region is expected to cause higher weathering rates and thus higher DIC in rivers (e.g., Gupta et al., 2011), but lower DIC concentrations were found in rivers of this region. It could be due to the influence of dilution because the dense precipitation over the small catchment area (Table 1) might have diluted DIC concentrations in rivers of this region. In order to understand the influence of the density of rainfall on DIC in rivers, we normalized the volume of discharge from the river with its catchment area. The catchment area normalized volume of discharge was found to be much higher in rivers from the SW region ( $1.71 \text{ m}^3 \text{ m}^{-2}$ ) than the rivers from SE ( $0.17 \text{ m}^3 \text{ m}^{-2}$ ), NE ( $0.6 \text{ m}^3 \text{ m}^{-2}$ ) and NW ( $0.32 \text{ m}^3 \text{ m}^{-2}$ ) regions of India. About three times higher catchment area normalized discharge might have diluted DIC concentrations in the rivers of the former region. A strong exponential decrease in DIC concentrations with increasing rainfall over the catchment ( $r^2= 0.72$ ,  $p<0.001$ ; Fig. 4a) also suggests that DIC concentration in the Indian rivers are strongly influenced by density of precipitation over the catchment.

Rivers of the SW region are relatively small in size, both in terms of catchment area (total catchment area:  $20 \times 10^3 \text{ km}^2$ ) and the length of the river (mean length: 126 km), than the rivers from other regions (SE, NE and NW) of India (Table 1). Since the contribution of DIC from in-stream processes, such as decomposition of organic matter, has been demonstrated to increase along the course of the fluvial network (Hotchkiss et al., 2015), possibly due to increase in the residence time of water (Catalan et al., 2016), the lowest DIC concentrations found in rivers from the SW region may also, at least partly, be due to their small size. Fairly good positive correlation between DIC concentrations and length of the rivers ( $r^2=0.38$ ,  $p<0.01$ ; Fig. 4b) also support this argument.

The major physical and biological processes controlling DIC concentrations in rivers are the exchange of  $\text{CO}_2$  with the atmosphere, autotrophic removal and heterotrophic addition of DIC. Since the Indian monsoonal estuaries have been reported to be a source of  $\text{CO}_2$  to the atmosphere during the discharge period due to heterotrophic decomposition of organic matter (Sarma et al., 2001, 2011, 2012; Gupta et al., 2008, 2009; Bhavya et al., 2018), the DIC input from the dissolution of atmospheric  $\text{CO}_2$  may be unlikely. On the other hand, organic matter decomposition is expected to add significant amount of DIC as

enhanced bacterial respiration rates were reported during this period (Sarma et al., 2011; 2012). In contrast, significant negative correlation between chlorophyll-*a* and DIC ( $r^2=-0.44$ ,  $p<0.01$ ; Fig. 4c), except few SE rivers where elevated phytoplankton biomass (Chl-*a*:  $>5 \text{ mg m}^{-3}$ ) was recorded, suggesting that autotrophic removal of DIC is also significant in the Indian monsoonal rivers during the study period. A significant positive relationship was observed between the  $\delta^{13}\text{C}_{\text{DIC}}$  and Chl-*a* ( $r^2=0.49$ ;  $p<0.01$ ; Fig. 4d), supporting this argument because preferential uptake of  $^{12}\text{C}$  than  $^{13}\text{C}$  during photosynthesis leaves the residual DIC enriched in  $^{13}\text{C}$ . On the other hand,  $\delta^{13}\text{C}_{\text{DIC}}$  showed significant positive correlation with DO saturation ( $r^2=0.50$ ,  $p<0.01$ ; Fig. 4e) (depleted  $\delta^{13}\text{C}_{\text{DIC}}$  values at more under saturation of DO) and DOC concentrations ( $r^2=0.43$ ,  $p<0.01$ ; Fig. 4f) as was observed in the Xi river (Zou et al., 2016). Altogether, enriched  $\delta^{13}\text{C}_{\text{DIC}}$  are associated with higher DOC, less under saturation of DO and higher phytoplankton biomass (Chl-*a*) while the depleted  $\delta^{13}\text{C}_{\text{DIC}}$  are associated more under saturation of DO and less DOC. This suggests that both autotrophic removal and heterotrophic addition control DIC in the Indian rivers during the discharge period, with a considerable spatial variability. However, influence of these processes on DIC concentrations is difficult to separate with this bulk  $\delta^{13}\text{C}_{\text{DIC}}$  data set, as the  $\delta^{13}\text{C}_{\text{DIC}}$  in rivers is also influenced by pollution, catchment lithology and outgassing of  $\text{CO}_2$  (Shin et al., 2011; Brunet et al., 2005; Bouillon et al., 2009; Zeng et al., 2011; Tamooh et al., 2013). Excluding Sabarmati and Mahisagar rivers, DIC concentrations showed fairly good linear relationship with population density over the catchment of the river ( $r^2=0.41$ ,  $p<0.01$ ; Fig. 4g), suggesting that considerable influence of pollution from the mega cities and industries on DIC in the Indian rivers.

Spatial distribution of soils shows that rivers of the NW region of India and upper reaches of Krishna and Godavari rivers drain the lime-rich black soils (Fig. 1) while rivers from the SW region drain red loamy soils. Whereas, the east-flowing rivers drain the lime-poor red sandy soils in the upper but lime-rich alluvial soils in the lower reaches (Fig.1). Lateritic soils, which are poor in lime and silicate, occupied the catchment of the rivers in the SW region of India. Relatively lower chemical weathering rates of the lateritic than the non-lateritic soils could be one of the reasons for the observed lower DIC concentration the rivers from SW region of India. A significant positive correlation was found between total alkalinity (TA) and  $\delta^{13}\text{C}_{\text{DIC}}$  ( $r^2=0.52$ ;  $p<0.01$ ; Fig. 4h), suggesting that significant contribution of DIC is from weathering of carbonate minerals in the catchment. Though the higher chemical weathering rates were reported for the Deccan Trap basalts (Das et al., 2005;

Singh et al., 2005), which occupied the catchments of rivers of the NW region of India and upper reaches of Godavari and Krishna, higher DIC concentrations were also observed in rivers draining over the metamorphic rocks. This suggests that the influences of factors other than bedrock are also significant on the concentrations of DIC in the Indian rivers.

#### 4.2 Major sources of DIC in the Indian monsoonal rivers

Though, the  $\delta^{13}\text{C}_{\text{DIC}}$  is a promising tool to decipher the sources of DIC, its interpretation for source material identification in rivers is still challenging because multiple physical and biological processes within the rivers significantly alter the  $\delta^{13}\text{C}$  of DIC source. The influence of major in-stream processes on the  $\delta^{13}\text{C}_{\text{DIC}}$  must be separated before interpreting the results for major sources of DIC, failing which leads to erroneous conclusions. In order to identify and separate DIC sources, we used here two different graphical mixing model techniques, Keeling plot (Keeling, 1958; Pataki et al., 2003) and Miller-Tans plots (Miller and Tans, 2003). These models approximate the hypothetical  $\delta^{13}\text{C}$  of source material as an intercept (in Keeling plot) and slope (in Miller-Tans plot) of the least square linear regression equations (Pataki et al., 2003; Campeau et al., 2017). The deviations from the approximated  $\delta^{13}\text{C}$  of source can be interpreted as the influence of in-stream processes. Further, we approximated the  $\delta^{13}\text{C}$  of  $\text{CO}_2$  using a set of enrichment factors of isotopic fractionation across the carbonate species (Zhang et al., 1995) in order to filter the impact of DIC speciation and pH on the bulk  $\delta^{13}\text{C}_{\text{DIC}}$  values. This approach has already been used by Quay et al. (1992), Mayorga et al. (2005) and recently by Campeau et al. (2017).

Significant negative relationships were observed in both Keeling plot ( $\delta^{13}\text{C}_{\text{DIC}}$  as a function of  $1/\text{DIC}$ ; Fig. 5a) and Miller-Tans plot ( $\delta^{13}\text{C}_{\text{DIC}} \times \text{DIC}$  as a function of  $\text{DIC}$ ; Fig. 5b) ( $r^2=0.61$ ,  $p<0.01$  and  $r^2=0.72$ ,  $p<0.01$  respectively) of DIC in the Indian rivers, except the rivers draining the Deccan Trap basalts. Both graphical mixing models, Keeling and Muller-Tans plots, approximated the similar  $\delta^{13}\text{C}$  of source material ( $-3.0\text{‰}$  and  $-2.0\text{‰}$  respectively; Fig. 5a&b), suggesting that weathering of carbonate minerals is the predominant source of DIC in the Indian monsoonal rivers rather than biogenic soil  $\text{CO}_2$ . Calculated  $\delta^{13}\text{C}$  of  $\text{CO}_2$  ranged from  $-21.5$  to  $-9.6\text{‰}$  in the Indian rivers with a mean value of  $-13.0\pm 2.7\text{‰}$ . Calculated  $\delta^{13}\text{C}$  of  $\text{CO}_2$  is linearly correlated with the measured  $\delta^{13}\text{C}_{\text{DIC}}$ , but correlation coefficient ( $r^2$ ) is only 0.51 (Fig. 5c), suggesting that significant spatial variability in the influence of in-stream processes on the  $\delta^{13}\text{C}_{\text{DIC}}$ . The Miller-Tans plot of  $\text{CO}_2$  ( $\delta^{13}\text{C}\text{-CO}_2 \times \text{CO}_2$  as a function of

CO<sub>2</sub>) showed highly significant linear regression model with a slope of -10.7‰ ( $r^2=0.97$ ;  $p<0.001$ ; Fig. 5d). These results indicated that chemical weathering of carbonate and silicate minerals by soil CO<sub>2</sub> (-10 to -9‰) is the major source of DIC in the Indian rivers. contributed by. Deviations of the measured  $\delta^{13}\text{C}_{\text{DIC}}$  (-13.0 to -1.4‰) from that of the approximated  $\delta^{13}\text{C}$  of DIC source (-3.0 to -2.0‰) and  $\delta^{13}\text{C}$  of CO<sub>2</sub> (-10.7‰) could be due to the influence of in-stream process. In more than 75% of the Indian rivers sampled, the deviation from the  $\delta^{13}\text{C}$  of DIC source is towards negative side (depletion) ( $\delta^{13}\text{C}_{\text{DIC}} < -3.0‰$ ), suggesting that heterotrophic decomposition of organic matter is the dominant process controlling DIC in these rivers. While, no (or very little) deviation was observed only in rivers from the SE region of India (mean  $\delta^{13}\text{C}_{\text{DIC}}$ : -3.1‰) could be due to the competition between autotrophy, degassing and heterotrophy as these processes influences the  $\delta^{13}\text{C}_{\text{DIC}}$  in opposite directions (Fig. 2); the former two processes causes enrichment while the latter depletes  $\delta^{13}\text{C}_{\text{DIC}}$ . Relatively higher phytoplankton biomass (mean Chl-a: 4.6 mg m<sup>-3</sup>) and less under-saturation of DO (98.7%) was observed in these rivers compared to the mean of the rest of the Indian rivers (2.4 mg m<sup>-3</sup> and 87.5% respectively), suggesting that autotrophy is one of the dominant processes controlling DIC in rivers from the SE region of India. Total number of dams on the rivers from this (SE) region (mean 155, Table 1) is not significantly higher from that of the mean of total number of dams on the Indian rivers sampled (mean 135) , suggesting that degassing due to storage of water may not be the dominant process responsible for enrichment in  $\delta^{13}\text{C}_{\text{DIC}}$  of these rivers.

### 4.3 Total DIC export by the Indian monsoonal rivers to the north Indian Ocean

Indian monsoonal rivers annually export ~10.3 Tg of DIC to the north Indian Ocean. Nearly three fourth of this amount (7.8 Tg) reaches to the Bay of Bengal while the Arabian Sea receives only one fourth (2.5 Tg). This is consistent with the higher magnitude of freshwater discharge to the Bay of Bengal (378 km<sup>3</sup> yr<sup>-1</sup>) from the catchment area of about 970x10<sup>3</sup> km<sup>2</sup> than the Arabian Sea (122 km<sup>3</sup> yr<sup>-1</sup> from the catchment area of 244x10<sup>3</sup> km<sup>2</sup>). The total DIC export by the Indian monsoonal rivers (10.3 Tg yr<sup>-1</sup>) is lower than the DIC export by the American (61.4 Tg yr<sup>-1</sup>) and African (17.7 Tg yr<sup>-1</sup>) rivers and major rivers draining to the tropical Atlantic from South America and Africa (53 Tg yr<sup>-1</sup>, Araujo et al. 2014). It is mainly due to the fact that freshwater discharge from the Indian monsoonal rivers is very low (~500 km<sup>3</sup> yr<sup>-1</sup>) compared to the American (11,799 km<sup>3</sup> yr<sup>-1</sup>) and African (3,786 km<sup>3</sup> yr<sup>-1</sup>) rivers. However, the Indian monsoonal rivers are exporting disproportionately

higher DIC because they account for only 1.3% of the global river discharge but export 2.5% of the global riverine DIC export to the oceans (400 Tg yr<sup>-1</sup>). Though American and African rivers account for 30% and 10% of the global river discharge, they export only 15% and 4.4% of global riverine DIC to oceans, respectively. Disproportionately higher DIC flux from the Indian rivers could be due to relatively higher weathering rates of silicate and carbonate minerals in their drainage basins (Das et al., 2005; Gurumurthy et al., 2012; Pattanaik et al., 2013). Higher DIC fluxes from the tropical regions are mainly attributed to the favourable climatic conditions, lithology and land use cover (Huang et al., 2012) of this region for higher dissolution.

Krishna et al. (2015) reported that Indian monsoonal rivers export 2.32 Tg yr<sup>-1</sup> of dissolved organic carbon (DOC) to the north Indian Ocean. The total fluvial dissolved carbon flux (DIC+DOC) would be 12.6 Tg yr<sup>-1</sup> in which DIC flux contributed up to ~81%. The predominance of DIC has also been found in rivers elsewhere in the world, for example, the British rivers (Jarvie et al., 2017) and high altitude Swedish rivers (Campeau et al., 2017). Since the catchment area of the Indian monsoonal rivers ranged widely from as low as 1x10<sup>3</sup> km<sup>2</sup> to as high as 313x10<sup>3</sup> km<sup>2</sup>, the export fluxes of DIC were normalized with the catchment area of the river (yield) in order to examine various factors controlling the DIC export to the north Indian Ocean.

#### **4.4 Yield of DIC from the Indian monsoonal rivers**

The yield of DIC found in this study (mean 8.7±5.2 g m<sup>-2</sup> yr<sup>-1</sup>) is close to those found in rivers from the tropical region of Asia, but significantly higher than those reported from tropical region of the American and African continents (Huang et al., 2012). The yield was highest (8.8±5.6 g m<sup>-2</sup> yr<sup>-1</sup>) in rivers from the SW region of India, despite they export relatively lower DIC (0.3 Tg yr<sup>-1</sup>) due to their low volume of discharge (46 km<sup>3</sup> yr<sup>-1</sup>) and relatively smaller catchment (20x10<sup>3</sup> km<sup>2</sup>) than the rivers from SE, NE and NW regions of India (Table 1. DIC yield showed a significant positive correlation with the volume of discharge ( $r^2=0.67$ ,  $p<0.001$ ; Fig. 6a) in medium rivers and no such relationship was found in the major rivers. Significant negative relationship was observed between DIC yield and catchment area of river ( $r^2 = -0.49$ ,  $p<0.001$ ; Fig. 6b and  $r^2 = -0.43$ ,  $p<0.001$ ; Fig. 6c for medium and major rivers respectively), suggesting the smaller rivers export more DIC per unit area of catchment compared to the major river systems, and thus inclusion of DIC data from medium rivers in the world significantly alters the global estimations of DIC. A fairly good linear relationship between the yield of DIC and the intensity of precipitation ( $r^2=0.43$ ,

$p < 0.01$  Fig. 6d) was observed only in the rivers which receives  $>2000\text{mm}$  of annual mean precipitation. Higher precipitation over the catchment increases the yield of DIC because the dense precipitation enhances the extraction of DIC from soils and rocks in their catchment. Therefore, high precipitation ( $\sim 3000\text{ mm}$ ) over the small catchment ( $20 \times 10^3\text{ km}^2$ ) could have increased DIC yield from the rivers of SW region of India.

Sreenivas et al. (2016) and Krishwan et al. (2009) found that the soil organic and inorganic carbon contents in the surface (100cm) soils in the catchment of rivers in the SW region were higher and lower, respectively, than the catchments of the rivers from SE, SW and NE region of India. Decomposition of soil organic matter releases excess  $\text{CO}_2$  and, the increase in soil  $\text{CO}_2$  leads to the formation of acidic conditions in soils. This would increase the DIC yield by more dissolution of soil carbonates and chemical weathering of carbonate and silicate rocks (Zou et al., 2016). A significant linear correlation was found between soil organic carbon content and DIC yield in this study ( $r^2=0.65$ ,  $p < 0.001$ ; Fig. 6e), suggesting that higher soil organic carbon in the catchment of the rivers from SW region could have elevated the yield of DIC from rivers of this region. The basin scale studies are, however, required for comprehensive understanding of the influence of environmental and anthropogenic factors on export fluxes and yield of DIC from the Indian monsoonal rivers.

## 5. Summary

In order to examine the spatial variability in the sources and distribution of dissolved inorganic carbon (DIC) in the Indian monsoonal rivers, and to estimate their export fluxes of DIC to the north Indian Ocean, we sampled a total of 27 major and medium rivers during wet period. An order of magnitude variability was found in DIC concentrations among the rivers sampled ( $3.4 - 73.6\text{ mg l}^{-1}$ ), with a lower mean concentration of  $6.6 \pm 2.1\text{ mg l}^{-1}$  in rivers located in the SW region of India. It is attributed to significant spatial variability in the size of rivers, precipitation pattern, pollution and lithology in their catchments. The approximated  $\delta^{13}\text{C}$  of DIC source from the Keeling and Miller-Tans plots ( $-2.0$  and  $-3.0\text{‰}$  respectively) and, the calculated  $\delta^{13}\text{C}$  of  $\text{CO}_2$  suggested that DIC in the Indian rivers is mainly originated from chemical weathering of carbonate minerals, but largely affected by autotrophic production in rivers from the southeast region of India and heterotrophic decomposition of organic matter in rivers from other regions of India. Indian monsoonal rivers together export  $\sim 10.3\text{ Tg yr}^{-1}$  of DIC to the north Indian Ocean, of which  $7.8\text{ Tg yr}^{-1}$  enters in to the Bay of Bengal while the Arabian Sea receives only  $2.5\text{ Tg yr}^{-1}$ . It is mainly attributed to the volume of river discharge as the former receives  $\sim 378\text{ km}^3\text{ yr}^{-1}$  while the latter receives only  $122\text{ km}^3$

yr<sup>-1</sup> from the Indian monsoonal rivers. Dense rainfall and higher soil organic carbon content in the catchment of rivers from the SW region than in the catchment of the other Indian rivers resulted in highest yield of DIC from the former than the latter.

## **6. Acknowledgements**

We would like to thank the Director, CSIR - National Institute of Oceanography (NIO), Goa, and the Scientist-In-Charge, NIO-Regional Centre, Visakhapatnam for their kind support and encouragement. We also acknowledge Dr. M. Dileep Kumar, NIO, Goa for his guidance and encouragement. The work is part of the Council of Scientific and Industrial Research (CSIR), funded research project. This publication has NIO contribution number .....

## **7. Data Availability**

The data set used in the current study can be obtained from the corresponding author by an e-mail request.

## **References**

Abril, G., Etcheber, H., Delille, B., Frankignoulle, M. and Borges A. V.: Carbonate dissolution in the turbid and eutrophic Loire estuary, *Mar. Ecol. Progr. Ser.*, 259, 129-138, 2003.

Abongwa, P. T., and Atekwana, E. A.: Assessing the temporal evolution of dissolved inorganic carbon in waters exposed to atmospheric CO<sub>2</sub>: a laboratory approach. *J. Hydrol.*, 505, 250-265, 2014.

Amiotte-Suchet, P. et al.:  $\delta^{13}\text{C}$  pattern of dissolved inorganic carbon in a small granitic catchment: the Strengbach case study (Vosges mountains, France), *Chem. Geol.*, 159, 129–145, doi:10.1016/s0009-2541(99)00037-6, 1999.

Araujo, M., Noriega, C., and Lefevre, N.: Nutrients and carbon fluxes in the estuaries of major rivers flowing into the tropical Atlantic, *Front. Mar. Sci.*, 1, 1-16, 2014.

Barber, R. T., Marra, J., Bidigare, R. C., Codispoti, L. A., Halpern, D., Johnson, Z., Latasa, M., Goericke, R., and Smith, S. L.: Primary productivity and its regulation in the Arabian Sea during 1995, *Deep Sea Res., Part II*, 48, 1127–1172, 2001.

Bauer, J. E., Cai, W. J., Raymond, P. A., Bianchi, T. S., Hopkinson, C. S., and Regnier, P.A.G.: The changing carbon cycle of the coastal ocean, *Nature*, 504, 61–70, doi:10.1038/Nature12857, 2013.

Berner, R. A., Lasaga, A. C., Garrels, R. M.: The carbonate–silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 million years, *Am. J. Sci.*, 283, 641-683, 1983

Bhavya, P. S., Sanjeev Kumar, Gupta, G. V. M., Sudharma, K. V., and Sudheesh, V.: Spatio-temporal variation in  $\delta^{13}\text{C}_{\text{DIC}}$  of a tropical eutrophic estuary (Cochin estuary, India) and adjacent Arabian Sea, *Continental Shelf Research*, 153, 75-85, doi:10.1016/j.csr.2017.12.006, 2018.

Boeglin, J. L., and Probst, J. L.: Physical and chemical weathering rates and  $\text{CO}_2$  consumption in a tropical lateritic environment: the upper Niger basin, *Chem. Geol.*, 148; 137-156, 1998.

Bouillon, S., Abril, G., Borges, A. V., Dehairs, F., Govers, G., Hughes, H. J., Merckx, R., Meysman, F. J. R., Nyunja, J., Osburn, C., and Middelburg, J. J.: Distribution, origin and cycling of carbon in the Tana River (Kenya): a dry season basin-scale survey from headwaters to the delta, *Biogeosciences*, 6, 2475–2493, doi:10.5194/bg-6-2475-2009, 2009.

Brunet, F. *et al.*  $\delta^{13}\text{C}$  tracing of dissolved inorganic carbon sources in Patagonian rivers (Argentina). *Hydrol. Process.*, 19, 3321–3344, doi:10.1002/hyp.5973, 2005.

Cai, W. J., Guo, X. H., Chen, C. T. A., Dai, M. H., Zhang, L. J., Zhai, W. D., Lohrenz, S. E., Yin, K. D., Harrison, P. J., and Wang, Y. C.: A comparative overview of weathering intensity and  $\text{HCO}_3^-$  flux in the world's major rivers with emphasis on the Changjiang, Huanghe, Zhujiang (Pearl) and Mississippi Rivers, *Cont. Shelf Res.*, 28, 1538-1549, 2008.

Cai, W. J.: Riverine inorganic carbon flux and rate of biological uptake in the Mississippi River plume, *Geophys. Res. Lett.*, 30, 1032, 2003.

Cai, W. –J.: Estuarine and coastal ocean carbon paradox:  $\text{CO}_2$  sinks or sites of terrestrial carbon incineration? *Annu Rev Mar Sci*, 3, 123-145, 2011.

Campeau, A., Wallin, M. B., Giesler, R., Löfgren, S., Mörth, C. –M., Schiff, S., Venkiteswaran, J. J. and Bishop, K.: Multiple sources and sinks of dissolved inorganic carbon across Swedish streams, refocusing the lens of stable C isotopes, *Nature, Scientific Reports*, 7, 9158, DOI:10.1038/s41598-017-09049-9, 2017.

Carritt, D. E. and Carpenter, J. H.: Comparison and evaluation of currently employed modifications of the Winkler method for determining dissolved oxygen in seawater: A NASCO report, *J. Mar. Res.*, 24, 286–318, 1966.

Catalan, N., Marce, R., Kothawala, D. N. & Tranvik, L. J. Organic carbon decomposition rates controlled by water retention time across inland waters. *Nature Geoscience* 9, 501–504, doi:10.1038/ngeo2720, 2016.

Central Water Commission, *Integrated Hydrological Data Book*, 680 pp., New Delhi, 2012.

Central Water Commission: *Integrated Hydrological Data Book*, 383 pp., New Delhi, 2006.

Christopher, P. B., Luettich Jr. R. A., Powers, S. P., Peterson, C. H., and McNinch, J. E.: Estimating the spatial extent of bottom-water hypoxia and habitat degradation in a shallow estuary, *Mar. Ecol. Prog. Ser.*, 230, 103–112, 2002.

Cole, J. J., Prairie, Y. T., Caraco, N. F., McDowell, W. H., Tranvik, L. J., Striegl, R. G., Duarte, C. M., Kortelainen, P., Downing, J. A., Middelburg, J. J., and Melack, J.: Plumbing



the global carbon cycle: Integrating inland waters into the terrestrial carbon budget, *Ecosystems*, 10, 171–184, 2007.

Coplen, T. B.: New guidelines for reporting stable hydrogen, carbon and oxygen isotope-ratio data, *Geochim. Cosmochim. Acta*, 60, 3359–3360, 1996.

Coplen, T. B. et al.: Compilation of minimum and maximum isotope ratios of selected elements in naturally occurring terrestrial materials and reagents, U.S. Department of the Interior and U.S. Geological Survey, 2002.

Das, A., Krishnaswami, S. and Bhattacharya, S. K.: Carbon isotope ratio of dissolved inorganic carbon (DIC) in rivers draining the Deccan Traps, India: Sources of DIC and their magnitudes. *Earth Planet. Sci. Lett.*, 236, 419–429, doi:10.1016/j.epsl.2005.05.009, 2005.

Deines, P., Langmuir, D., and Harmon, R. S.: Stable carbon isotope ratios and the existence of a gas phase in the evolution of carbonate ground waters, *Geochim. Cosmochim. Acta*, 38, 1147–1164, doi:10.1016/0016-7037(74)90010-6, 1974.

Dessert, C., Dupre, B., Francois, L. M., Schott, J., Gaillardet, J., Chakrapani, G., and Bajpai, S.: Erosion of Deccan Traps determined by river geochemistry: impact on the global climate and the  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of seawater. *Earth and Planet. Sci. Lett.*, 188, 459–474, 2001.

Dhillon, G. S., and Inamdar, S.: Extreme storms and changes in particulate and dissolved organic carbon in runoff: Entering uncharted waters?, *Geophys. Res. Lett.*, 40, 1322–1327, doi:10.1002/grl.50306., 2013.

Deirmendjian, L., and Abril, G.: Carbon dioxide degassing at the groundwater-stream atmosphere interface: isotopic equilibration and hydrological mass balance in a sandy watershed, *J. Hydrol.*, 2018, doi: <https://doi.org/10.1016/j.jhydrol.2018.01.003>

Findlay, S.: Stream microbial ecology, *J. North Am. Benthol. Soc.*, 29, 170–181, doi:10.1899/09-023.1, 2010.

Finlay, J. C. and Kendall, C.: Stable isotope tracing of temporal and spatial variability in organic matter sources to freshwater ecosystems, In: *Stable Isotopes in Ecology and Environmental Science*, 2nd edn., edited by: Michener, R. H. and Lajtha, K., Blackwell Publishing, Malden, USA, 283–333, 2007.

Finlay, J. C.: Controls of stream water dissolved inorganic carbon dynamics in a forested watershed, *Biogeochem.*, 62, 231–252, 2003.

Finlay, J. C.: Patterns and controls of lotic algal stable carbon isotope ratios, *Limnol. Oceanogr.*, 49, 850–861, 2004.

Gaillardet, J., Dupre, B., Louvat, P., and Allegre, C. J.: Global silicate weathering and  $\text{CO}_2$  consumption rates deduced from the chemistry of large rivers. *Chem. Geol.* 159, 3–30, 1999.

Gauns, M., Madhupratap, M., Ramaiah, N., Jyothibabu, R., Fernandes, V., Paul, J. T., and Kumar, S. P.: Comparative accounts of biological productivity characteristics and estimates of carbon fluxes in the Arabian Sea and the Bay of Bengal, *Deep Sea Res., Part II*, 52, 2003–2017, 2005.

- Garcia, E. H., and Gordon, L. I.: Oxygen solubility in seawater better fitting equations, *Limnol. Oceanogr.*, 37, 1307–1312, doi:10.4319/lo.1992.37.6.1307, 1992.
- Giesler, R. *et al.* Spatiotemporal variations of  $p\text{CO}_2$  and  $\delta^{13}\text{C}\text{-DIC}$  in subarctic streams in northern Sweden. *Global Biogeochemical Cycles* 27, 176–186, doi:10.1002/gbc.20024, 2013.
- Gupta, G.V.M., Sarma, V.V.S.S., Robin, R.S., Raman, A.V., Jai Kumar, M., Rakesh, M. and Subramanian, B. R.: Influence of net ecosystem metabolism in transferring riverine organic carbon to atmospheric  $\text{CO}_2$  in a tropical coastal lagoon (Chilka Lake, India). *Biogeochemistry*, 87, 265-285, doi:10.1007/s10533-008-9183-x, 2008.
- Gupta, G. V. M., Thottathil, S. D., Balachandran, K. K., Madhu, N. V., Madeswaran, P., and Nair, S.:  $\text{CO}_2$  supersaturation and net heterotrophy in a tropical estuary (Cochin, India): influence of anthropogenic effect, *Ecosystems*, 12, 1145–1157, doi:10.1007/s10021-009-9280-2, 2009.
- Gupta, H., Chakrapani, G. J., Selvaraj, K., and Kao, S.-J.: The fluvial geochemistry, contributions of silicate, carbonate and saline–alkaline components to chemical weathering flux and controlling parameters: Narmada River (Deccan Traps), India, *Geochim. Cosmochim. Acta*, 75, 800-824, 2011.
- Gurumurthy G. P., Balakrishna K., Riotte J., Braun J.-J., Audry S., Shankar H. N. U. and Manjunatha B. R.: Controls on intense silicate weathering in a tropical river, southwestern India, *Chem. Geol.*, 300–301, 61–69, 2012.
- Hotchkiss, E. R. *et al.* Sources of and processes controlling  $\text{CO}_2$  emissions change with the size of streams and rivers, *Nature Geoscience* 8, doi:10.1038/Ngeo2507, 2015.
- Huang, T-H., Fu, Y-H., Pan, P-Y., and Arthur, C.T.: Fluvial carbon fluxes in tropical rivers, *Current Opinion in Environmental Sustainability*, 4, 162–169, 2012.
- Ishikawa, N.F., Tayasu, I., Yamane, M., Yokoyama, Y., Sakai, S., and Ohkouchi, N.: Sources of dissolved inorganic carbon in two small streams with different bedrock geology; Insights from carbon isotopes. *Radiocarbon*, 57, 439–448, 2015.
- Jarvie, H.P., King, S.M., and Neal, C.: Inorganic carbon dominates total dissolved carbon concentrations and fluxes in British rivers: Application of the THINCARB model – Thermodynamic modeling of inorganic carbon in freshwaters, *Sci. Tot. Environ.*, 575, 496-512, 2017.
- Joesoef, A., Kirchman, D. L., Sommerfield, C.K., and Cai, W-J.: Seasonal variability of the inorganic carbon system in a large coastal plain estuary. *Biogeosciences*, 14, 4949-4963, 2017
- Keeling, C. D.: The concentration and isotopic abundances of atmospheric carbon dioxide in rural areas. *Geochimica et Cosmochimica Acta* 13, 322–334, doi:10.1016/0016-7037(58)90033-4 (1958).
- Kishwan, J., Pandey, R., and Dhadwal, V. K.: India's forest and tree cover: Contribution as a carbon sink, Tech. Pap. 130, ICFRE BL-23, 2009.

- Krishna, M. S., Prasad, V. R., Sarma, V. V. S. S., Reddy, N. P. C., Hemalatha, K. P. J., and Rao Y. V.: Fluxes of dissolved organic carbon and nitrogen to the northern Indian Ocean from the Indian monsoonal rivers, *J. Geophys. Res. Biogeosci.*, 120, 2067–2080, 2015.
- Kumar, R., Singh, R. D., and Sharma, K. D.: Water resources of India, *Curr. Sci.*, 89, 794–811, 2005.
- Land, L. S.: The isotopic and trace element geochemistry of dolomite: the state of the art. *Concepts and Models of Dolomitization*, 63, 485, doi:10.2110/pec.80.28.0087, 1980.
- Lerman, A., Wu, L. L., and Mackenzie, F. T.: CO<sub>2</sub> and H<sub>2</sub>SO<sub>4</sub> consumption in weathering and material transport to the ocean, and their role in the global carbon balance, *Mar. Chem.*, 106, 326-350, 2007.
- Löfgren, S., Froberg, M., Yu, J., Nisell, J. & Ranneby, B.: Water chemistry in 179 randomly selected Swedish headwater streams related to forest production, clear-felling and climate. *Environ Monit Assess* 186, 8907–8928, doi:10.1007/s10661-014-4054-5, 2014.
- Ludwig, W., Amiotte-Suchet, P., Munhoven, G., and Probst, J. L.: Atmospheric CO<sub>2</sub> consumption by continental erosion: present-day controls and implications for the last glacial maximum. *Global Planet Change*, 17, 107-120, 1998.
- Mackenzie, F. T., Lerman, A., and Andersson, A.J.: Past and present of sediment and carbon biogeochemical cycling models. *Biogeosciences* 1, 11 -32, 2004.
- Madhupratap, M., Prasanna Kumar, S., Bhattathiri, P. M. A., Kumar, M. D., Raghukumar, S., Nair, K. K. C., and Ramaiah, N.: Mechanism of the biological response to winter cooling in the northeastern Arabian Sea, *Nature*, 384, 549–552, 1996.
- Maher, D. T., Cowley, K., Santos, I. R., Macklin, P., and Eyre, B. D.: Methane and carbon dioxide dynamics in a subtropical estuary over a diel cycle: Insights from automated *in situ* radioactive and stable isotope measurements, *Mar. Chem.*, 168, 69–79, doi:10.1016/j.marchem.2014.10.017, 2015.
- Maher, D. T., Santos, I. R., Golsby-Smith, L., Gleeson, J., and Eyre, B. D.: Groundwater-derived dissolved inorganic and organic carbon exports from a mangrove tidal creek: The missing mangrove carbon sink?, *Limnol. Oceanogr.*, 58, 475-488, 10.4319/lo.2013.58.2.0475, 2013.
- Mayorga, E. *et al.* Young organic matter as a source of carbon dioxide outgassing from Amazonian rivers. *Nature* **436**, 538–541, doi:10.1038/nature03880, 2005.
- McConnaughey, T. A., LaBaugh, J. W., Rosenberry, D. O., Striegl, R. G., Reddy, M. M., Schuster, P. F., and Carter, V.: Carbon budget for a groundwater-fed lake: calcification supports summer photosynthesis, *Limnol. Oceanogr.*, 39, 1319–1332, 1994.
- Meybeck, M., and Ragu, A.: GEMS/water contribution to the Global Register of River Inputs (GLORI), Provisional Final Rep., 245 pp., UNEP/WHO/UNESCO, Geneva, Switzerland, 1995.
- Meybeck, M., and Ragu, A.: River discharges to the oceans. An assessment of suspended solids, major ions, and nutrients, *Environ. Inf. and Assess. Rep.*, 250, 1996.

Meybeck, M., and Vorosmarty, C. J.: Global transfer of carbon by rivers, *Global Change News Lett*, 37, 41974, 1999.

Meybeck, M.: Global chemical weathering of surficial rocks estimated from river dissolved loads, *Am. J. Sci.*, 287, 401–428, 1987.

Meybeck, M.: Riverine Transport of atmospheric carbon-sources, global typology and budget. *Water Air Soil Pollut.*, 70, 443-463, 1993.

Miller, J. B. & Tans, P. P.: Calculating isotopic fractionation from atmospheric measurements at various scales. *Tellus B* 55, 207–214, doi:[10.1034/j.1600-0889.2003.00020.x](https://doi.org/10.1034/j.1600-0889.2003.00020.x) (2003).

Mukhopadhyay, S. K., Biswas, H., De, T. K., Sen, S., and Jana, T. K.: Seasonal effects on the air–water carbon dioxide exchange in the Hooghly estuary, NE coast of Bay of Bengal, India. *J Environ Monit.*, 4, 549-552, 2002.

Muraleedharan, P. M., and Prasanna Kumar, S.: Arabian Sea upwelling—A comparison between coastal and open ocean regions, *Curr. Sci.*, 71, 842–846, 1996.

O’Leary, M. H.: Carbon Isotopes in Photosynthesis, *BioScience*, 38, 328–336, doi:[10.2307/1310735](https://doi.org/10.2307/1310735), 1988.

Opsahl, S. P. and Zepp, R. G.: Photochemically-induced alteration of stable carbon isotope ratios ( $\delta^{13}\text{C}$ ) in terrigenous dissolved organic carbon, *Geophys. Res. Lett.*, 28, 2417–2420, doi:[10.1029/2000gl012686](https://doi.org/10.1029/2000gl012686), 2001.

Parker, S. R., Poulson, S. R., Gammons, C. H., and DeGrandpre, M. D.: Biogeochemical controls on diel cycling of stable isotopes of dissolved  $\text{O}_2$  and dissolved inorganic carbon in the Big Hole River, Montana, *Environ. Sci. Tech.*, 39, 7134–7140, doi:[10.1021/es0505595](https://doi.org/10.1021/es0505595), 2005.

Parker, S. R., Poulson, S. R., Smith, M. G., Weyer, C. L., and Bates, K. M.: Temporal variability in the concentration and stable carbon isotope composition of dissolved inorganic and organic carbon in two Montana, USA Rivers, *Aquat Geochem.*, 16, 61–84, doi:[10.1007/s10498-009-9068-1](https://doi.org/10.1007/s10498-009-9068-1), 2010.

Pataki, D. E. *et al.* The application and interpretation of Keeling plots in terrestrial carbon cycle research. *Global Biogeochemical Cycles* 17, 1022, doi:[10.1029/2001GB001850](https://doi.org/10.1029/2001GB001850) (2003).

Pattanaik, J. K., Balakrishnan, S., Bhutani, R. and Singh, P.: Estimation of weathering rates and  $\text{CO}_2$  drawdown based on solute load: Significance of granulites and gneisses dominated weathering in the Kaveri River basin, Southern India, *Geochim. Cosmochim. Acta*, 121, 611-636, 2013.

Pattanaik, S., Sahoo, R. .K, Satapathy, D. R., Panda, C. R., Choudhury, S. B. et al.: Intra-annual Variability of  $\text{CO}_2$  Flux in the Mahanadi Estuary- A Tropical Estuarine System, India. *Ann Mar Sci.*, 1, 005-012, 2017.

Polsenaere, P., and Abril, G.: Modelling  $\text{CO}_2$  degassing from small acidic rivers using water  $\text{pCO}_2$ , DIC and  $\delta^{13}\text{C}$ -DIC data. *Geochim. Cosmochim. Acta* 91, 220–239, 2012.

doi:[10.1016/j.gca.2012.05.030](https://doi.org/10.1016/j.gca.2012.05.030)

Prasanna Kumar, S., Muraleedharan, P. M., Prasad, T. G., Gauns, M., Ramaiah, N., de Souza, S. N., Sardesai, S., and Madhupratap, M.: Why is the Bay of Bengal less productive during summer monsoon compared to the Arabian Sea?, *Geophys. Res. Lett.*, 29, 2235, doi:10.1029/2002GL016013, 2002.

Prasanna Kumar, S., Nuncio, M., Narvekar, J., Kumar, A., Sardesai, S., DeSousa, S. N., Gauns, M., Ramaiah, N., and Madhupratap, M.: Are eddies nature's trigger to enhance biological productivity in the Bay of Bengal? *Geophys. Res. Lett.* 31, 5. doi:10.1029/2003G1019274, 2004.

Quay, P. D. *et al.* Carbon cycling in the Amazon River: Implications from the  $^{13}\text{C}$  compositions of particles and solutes. *Limnology and Oceanography* 37, 857–871, doi:10.4319/lo.1992.37.4.0857, 1992.

Raymond, P. A., and Cole, J. J.: Increase in the export of alkalinity from North America's largest river, *Science*, 301, 88–91, doi:10.1126/science.1083788, 2003.

Raymond, P. A., Oh, N. H., Turner, R. E., and Broussard, W.: Anthropogenically enhanced fluxes of water and carbon from the Mississippi River, *Nature*, 451, 449–452, doi:10.1038/Nature06505, 2008.

Raymond, P.A., and Bauer, J.: Atmospheric  $\text{CO}_2$  evasion, dissolved inorganic carbon production, and net heterotrophy in the York River estuary, *Limnol.Oceanogr.*, 45, 1707–1717, 2000.

Regnier, P., et al.: Anthropogenic perturbation of the carbon fluxes from land to ocean, *Nat. Geosci.*, 6, 597–607, doi:10.1038/ngeo1830, 2013.

Ren, W., Tian, H., Tao, B., Yang, J., Pan, S., Cai, W.-J., Lohrenz, S. E., He, R., and Hopkinson, C. S.: Large increase in dissolved inorganic carbon flux from the Mississippi River to Gulf of Mexico due to climatic and anthropogenic changes over the 21st century, *J. Geophys. Res. Biogeosci.*, 120, 724–736, doi:10.1002/2014JG002761, 2015.

Rengarajan, R., and Sarma, V. V. S. S.: Submarine groundwater discharge and nutrient addition to the coastal zone of the Godavari estuary, *Mar. Chem.*, 172, 57–69, 2015.

Richey, J.E. et al: Outgassing from Amazonian rivers and wetlands as a large tropical source of atmospheric  $\text{CO}_2$ , *Nature* 416, 617–620, 2002

Samanta, S., Dalai, T. K., Pattanai K. J. K., Rai, S. K., and Mazumdar, A.: Dissolved inorganic carbon (DIC) and its  $\delta^{13}\text{C}$  in the Ganga (Hooghly) River estuary, India: Evidence of DIC generation via organic carbon degradation and carbonate dissolution. *Geochim.Cosmochim.Acta*, 165, 226–248, 2015.

Sarma, V. V. S. S., et al.: Emission of carbon dioxide from the Indian monsoonal estuaries, *Geophys. Res. Lett.*, 39, L03602, doi:10.1029/2011GL050709, 2012.

Sarma, V. V. S. S., et al.: High  $\text{CO}_2$  emissions from the tropical Godavari estuary (India) associated with monsoon river discharges, *Geophys. Res. Lett.*, 38, L08601, doi:10.1029/2011GL046928, 2011.

- Sarma, V. V. S. S., Kumar, M. D., and Manerikar, M.: Emission of carbon dioxide from a tropical estuarine system, Goa, India, *Geophys. Res. Lett.*, 28, 1239–1242, doi:10.1029/2000GL006114, 2001.
- Sarma, V. V. S. S., Krishna, M. S., Prasad, V. R., Kumar, B. S. K., Naidu, S. A., Rao, G. D., Viswanadham, R., Sridevi, T., Kumar, P. P., and Reddy, N. P. C.: Distribution and sources of particulate organic matter in the Indian monsoonal estuaries during monsoon, *J. Geophys. Res. Biogeosci.*, 119, doi:10.1002/2014JG002721, 2014.
- Shetye, S. R., Gouveia, A. D., Shenoi, S. S. C.: Circulation and water masses of the Arabian Sea, *Proc. Indian Acad. Sci. Earth Planet. Sci.*, 103, 107–123, 1994.
- Shin, W. J., Chung, G. S., Lee, D., and Lee, K. S.: Dissolved inorganic carbon export from carbonate and silicate catchments estimated from carbonate chemistry and  $\delta^{13}\text{C}_{\text{DIC}}$ . *Hydrol. Earth Syst. Sci.*, 15, 2551 – 2560, 2011.
- Singh, S.K., Sarin, M.M., and France-Lanord, C.: Chemical erosion in the eastern Himalaya: Major ion composition of the Brahmaputra and  $^{13}\text{C}$  of dissolved inorganic carbon. *Geochim. Cosmochim. Acta*, 69, 3573-3588, 2005.
- Smith S. L.: Understanding the Arabian Sea: reflections on the 1994–1996 Arabian Sea expedition, *Deep Sea Res. Part II* 48, 1385–1402, 2001.
- Solomon, D. K., and Cerling, T. E.: The annual carbon dioxide cycle in a montane soil: observations, modeling and implications for weathering, *Water Resources Res.*, 23, 2257-2265, 1987.
- Soman, M. K., and Kumar, K. K.: Some aspects of daily rainfall distribution over India during the southwest monsoon season, *Int. J. Clim.*, 19, 299–311, 1990.
- Sreenivas, K., Dadhwal, V. K., Suresh, K., Sri Harsha, G., Tarik, M., Sujatha, G, Suresh, J. R., G., Fyzee, M., and Ravisankar, T.: Digital mapping of soil organic and inorganic carbon status in Indi, *Geoderm.*, 269. 160-173, 10.1016/j.geoderma.2016.02.002, 2016.
- Sridevi, B., Sarma, V.V.S.S., Murty, T.V.R., Sadharam, Y., Reddy, N.P.C., Vijayakumar, K., Raju, N.S.N., Jawahar Kumar, Ch., Raju, Y.S.N., Luis, R., Kumar, M.D., Prasad, K.V.S.R. : Variability in stratification and flushing times of the Gautami–Godavari estuary, India, *J. Earth. Sys., Sci.*, 124, 993-1003, 2015.
- Subramanian, V.: Sediment load of Indian rivers, *Curr. Sci.*, 64, 928–930, 1993.
- Suzuki, R., and Ishimaru, T.: An improved method for the determination of phytoplankton chlorophyll using N,N-dimethyl formamide, *J. Oceanogr.*, 46, 190–194, 1990.
- Tamooh, F., Borges, A. V., Meysman, F. J. R., Meersche, K. V. D., Dehairs, F., Merckx, R. et al.: Dynamics of dissolved inorganic carbon and aquatic metabolism in the Tana River Basin, Kenya, *Biogeosciences Discuss.*, 10, 5175–5221, 2013.
- Tamse, S., Ogrinc, N., Walter, L.M, Turk, D., and Faganeli, J.: River Sources of Dissolved Inorganic Carbon in the Gulf of Trieste (N Adriatic): Stable Carbon Isotope Evidence, *Estuaries and Coasts*, DOI 10.1007/s12237-014-9812-7, 2014.

- Tank, J. L., Rosi-Marshall, E. J., Griffiths, N. A., Entrekin, S. A., and Stephen, M. L.: A review of allochthonous organic matter dynamics and metabolism in streams, *J. North Am. Benthol. Soc.*, 29, 118–146, doi:10.1899/08-170.1, 2010.
- Vähätalo, A. V., and Wetzel, R. G.: Long-term photochemical and microbial decomposition of wetland-derived dissolved organic matter with alteration of  $^{13}\text{C}:^{12}\text{C}$  mass ratio, *Limnol. Oceanogr.*, 53, 1387–1392, doi:10.4319/lo.2008.53.4.1387, 2008.
- Varkey, M. J., Murty, V. S. N., and Suryanarayana, A.: Physical oceanography of the Bay of Bengal and Andaman Sea, *Oceanogr. Mar. Biol.*, 34, 1–70, 1996.
- Venkiteswaran, J.J., Schiff, S.L., and Wallin, M.B.: Large Carbon Dioxide Fluxes from Headwater Boreal and Sub-Boreal Streams. *PLoS ONE* 9, e101756, 2014.
- Viers, J., Oliva, P., Dandurand, J. L., Dupré, B., Gaillardet, J., Heinrich, D. H., and Karl, K. T.: Chemical weathering rates,  $\text{CO}_2$  consumption, and control parameters deduced from the chemical composition of Rivers, *Treatise on Geochemistry* Pergamon, Oxford, 2007.
- Vijith, V., Sundar, D., and Shetye, S. R.: Time-dependence of salinity in monsoonal estuaries, *Estuar. Coast. Shelf Sci.*, 85, 601–608, doi:10.1016/j.ecss.2009.10.003, 2009.
- Waldron, S., Hall, A. J., and Fallick, A. E.: Enigmatic stable isotope dynamics of deep peat methane, *Glob. Biogeochem. Cy.*, 13, 93–100, doi:10.1029/1998gb900002, 1999.
- Waldron, S., Scott, E. M., and Soulsby, C.: Stable isotope analysis reveals lower-order river dissolved inorganic carbon pools are highly dynamic, *Environ. Sci. Technol.*, 41, 6156–6162, doi:10.1021/es0706089, 2007.
- Wang, Z. A., Bienvenu, D.J., Mann, P.J., Hoering, K.A. Poulsen, J.R., Spencer, R.G.M., and Holmes, R.M.: Inorganic carbon speciation and fluxes in the Congo River, *Geophys. Res. Lett.*, 40, 511–516, 2013.
- Williamson, C. E., Zagarese, H. E., Schulze, P. C., Hargreaves, B. R., and Seva, J.: The impact of short-term exposure to UV-B radiation on zooplankton communities in north temperate lakes, *J. Plankton Res.*, 16, 205–218, doi:10.1093/plankt/16.3.205, 1994.
- Zeng, F- W., Masiello C. A., and Hockaday, W. C.: Controls on the origin and cycling of riverine dissolved inorganic carbon in the Brazos River, Texas, *Biogeochemistry*, 104, 275–291, doi:10.1007/s10533-010-9501-y, 2011.
- Zou, J.: Sources and Dynamics of Inorganic Carbon within the Upper Reaches of the Xi River Basin, Southwest China, *PLoS One*, 11, e0160964. doi:10.1371/journal.pone.0160964, 2016.

## Figure captions

**Figure 1:** Map showing the study region. Rivers sampled in this study were indicated by solid black line. Distribution of soils in catchments of the Indian monsoonal rivers sampled was also shown. Rivers draining the four regions, i.e., northwest (NW), southwest (SW), southeast (SE) and northeast (NE) were shown by solid black arrows. Source: [https://www.clearias.com/up/Major\\_soil\\_types\\_in\\_India.jpg](https://www.clearias.com/up/Major_soil_types_in_India.jpg)

**Figure 2:** Schematic diagram showing the typical range of  $\delta^{13}\text{C}$  of different sources of dissolved inorganic carbon (DIC) in rivers. Various major processes influencing the  $\delta^{13}\text{C}$  of DIC ( $\delta^{13}\text{C}_{\text{DIC}}$ ) within the rivers were also shown. Hollow black arrows ( $\leftarrow$  and  $\rightarrow$ ) indicates the direction of change in  $\delta^{13}\text{C}_{\text{DIC}}$  due to the influences different in-stream process mentioned against arrows.

**Figure 3:** Spatial variability in the concentration ( $\text{mg l}^{-1}$ ; 3a), export flux ( $\text{Tg yr}^{-1}$ ; 3b) and yield ( $\text{g m}^{-2} \text{yr}^{-1}$ ; 3c) of dissolved inorganic carbon (DIC) and its stable isotopes ( $\delta^{13}\text{C}_{\text{DIC}}$ , 3d) in the Indian monsoonal rivers studied. Rivers geographically located in the northwest (NW), southwest (SW), southeast (SE) and northeast (NE) regions of India were also shown. Rivers draining into the Bay of Bengal (east-flowing rivers) were shown by gray shade while rivers draining into the Arabian Sea (west-flowing) were shown by no shade.

**Figure 4:** (a) Exponential decrease and (b) linear increase of dissolved inorganic carbon (DIC) concentrations with increasing the rainfall over the catchment and length of the river respectively. (c) Inverse and (d) linear relationships of chlorophyll-a (Chl-a) with concentrations and  $\delta^{13}\text{C}$  of DIC respectively. Significant linear relationships of  $\delta^{13}\text{C}$  of DIC with (e) dissolved oxygen (DO) saturation and (f) dissolved organic carbon (DOC) concentration. Linear relationships observed between (g) DIC concentrations and population density in the catchment and (h) total alkalinity and  $\delta^{13}\text{C}$  of DIC in the Indian monsoonal rivers during the study period. Ovals with dashed line indicate the outliers which were not included in the regression equations. Rivers of the northwest region of India showed linear relationships as shown by the other Indian rivers but with a different slope (Fig. f-h)

**Figure 5:** Least square linear regression models of (a)  $\delta^{13}\text{C}_{\text{DIC}}$  as a function of  $1/\text{DIC}$  (Keeling plot) and (b)  $\delta^{13}\text{C}_{\text{DIC}} \times \text{DIC}$  as a function of DIC concentrations (Miller-Tans plot) in the Indian monsoonal rivers. (c) Linear relationship between calculated  $\delta^{13}\text{C}$  of  $\text{CO}_2$  and the measured  $\delta^{13}\text{C}_{\text{DIC}}$  and (d) Miller-Tans linear regression model of  $\delta^{13}\text{C}-\text{CO}_2 \times \text{CO}_2$  as a function of  $\text{CO}_2$  concentration in the Indian monsoonal rivers.

**Figure 6:** Significant relationships of dissolved inorganic carbon (DIC) yield with (a) river discharge in medium estuaries, (b) catchment areas of the medium rivers, (c) catchment areas of the major rivers, (d) rainfall over the catchment of all the rivers sampled and (e) soil organic carbon (OC) content in catchments of the Indian monsoonal rivers studied. Since the data on soil OC is not available for each watershed (e) was plotted using the available soil OC data on regional scale (NW, SW, SE and NE regions of India), Hence, it contains only four points.




## Table captions



**Table 1:** Catchment area, discharge, length and elevation of each river and, annual mean rainfall, number of dams and population density in each watershed of the Indian monsoonal rivers sampled. Concentrations, export fluxes and yields of dissolved inorganic carbon (DIC) and its stable isotopes ( $\delta^{13}\text{C}_{\text{DIC}}$ ) of Indian rivers were given. Measured pH and calculated  $\delta^{13}\text{C}$  of  $\text{CO}_2$  from isotopic fractionation factors across DIC speciation were also provided. Rivers located in the northern region (north of  $16^\circ\text{N}$ ) of India were shown by the shaded (grey) area. Of these rivers, Mahisagar, Sabarmati, Tapti and Narmada are located in the northwestern (NW) whereas, rivers from Godavari to Hyadri are located in the northeastern (NE) region of India.

River name	Catchment area (x10 <sup>3</sup> km <sup>2</sup> )	Annual Discharge (km <sup>3</sup> )	Length (km)	Population (No. km <sup>-2</sup> )	No. of major Dams	Elevation (m)	Rain-fall (mm)	pH	DIC Conc. (mg l <sup>-1</sup> )	DIC export flux (Tg yr <sup>-1</sup> )	DIC yield (g m <sup>-2</sup> yr <sup>-1</sup> )	δ <sup>13</sup> C <sub>DIC</sub> (‰)	Cal. δ <sup>13</sup> C-CO <sub>2</sub> (‰)
<b>West flowing rivers (Arabian Sea)</b>													
MAHISAGAR	34.8	11.0	580	507	138	500	785	7.3	22.8	0.3	7.2	-12.4	-21.5
SABARMATI	21.7	3.8	371	1702	62	1173	787	7.2	24.6	0.1	4.3	-13.0	-
TAPTI	65	14.9	724	208	375	752	888	7.1	35.6	0.5	8.2	-7.9	-16.4
NARMADA	99	45.6	1312	184	281	1317	1120	7.5	30.6	1.4	14.1	-11.0	-
ZUARI	1	3.2	34	92	3	-	3500	7.0	4.8	0.0	4.4	-5.1	-13.2
MANDOVI	3.6	3.3	81	62	2	600	3500	6.5	4.5	0.0	14.5	-7.4	-13.6
KHALI	4.2	4.8	184	111	6	600	3200	7.2	5.7	0.0	6.5	-7.0	-15.5
SHARAVATHI	3.6	4.5	128	109	3	700	4000	6.5	8.4	0.0	10.6	-6.8	-13.1
NETRAVATHI	3.2	11.1	103	103	-	1000	3923	6.3	5.1	0.1	17.8	-7.1	-12.4
BHARATHAPUZHA	6.2	5.1	209	-	13	1964	2500	6.0	3.4	0.0	2.8	-7.9	-11.6
CHALAKUDY	1.7	1.9	144	-	6	1250	3600	6.3	4.8	0.0	5.4	-5.1	-10.3
<b>East flowing rivers (Bay of Bengal)</b>													
VAIGAI	7	1.1	258	499	2	1200	850	-	26.0	0.0	4.2	-7.9	-
AMBALAYAAR	-	0.9	-	-	-	-	-	8.6	32.2	0.0	-	-	-
CAUVERY	88	21.3	800	393	122	1341	1075	7.5	40.5	0.9	9.8	-2.9	-11.8
VELLAR	8.6	0.9	210	457	3	900	980	7.4	73.6	0.1	7.7	-2.4	-11.1
PONNAIYAR	16	1.6	396	291	4	900	969	7.4	43.4	0.1	4.3	-1.4	-10.3
PENNA	55	6.3	597	186	61	1439	510	8.3	37.1	0.2	4.3	-2.6	-11.9
KRISHNA	259	69.8	1300	260	736	1903	784	8	33.5	2.3	9.0	-3.9	-
GODAVARI	313	110.5	1465	193	978	1067	1300	-	13.1	1.5	4.6	-5.6	-9.6
NAGAVALI	9.4	2.0	256	150	4	1300	1000	-	15.5	0.0	3.3	-	-13.1
VAMSADHARA	11.0	3.5	254	130	3	370	1400	8.2	27.2	0.1	8.6	-3.0	-12.4
RUSIKULYA	9.0	1.9	175	360	13	1000	1000	-	19.8	0.0	4.2	-	-
MAHANADI	141.6	66.9	858	282	280	890	1406	6.9	13.3	0.9	6.3	-5.4	-13.2
BAITARANI	14.2	28.5	414	324	8	900	1450	6.0	10.3	0.3	20.7	-6.5	-10.2
SUBARNALEKHA	29.2	12.4	395	338	12	600	1800	8.0	13.6	0.2	5.8	-2.9	-12.0
HYADRI	10.2	50.5		191	-	-	-	7.1	24.1	1.2	-	-5.8	-14.0


**Table 1**




**Fig. 1**




**Fig. 2**




**Fig. 3**



**Fig. 4**



**Fig.5**



**Fig. 6:**