

## **Influence of hydrological factors on bacterial community structure in a tropical monsoonal estuary in India**

**Parvathi Ammini<sup>1\*</sup>, Michela R Catena<sup>1</sup>, Jasna Vijayan<sup>1,2</sup>, Nikhil Phadke<sup>3</sup>, Nikhita Gogate<sup>3</sup>**

<sup>1</sup>CSIR-National Institute of Oceanography, Regional Centre (CSIR), Kochi-682 018, India

<sup>2</sup>Department of Aquatic Life Medicine, College of Fisheries Science, Pukyong National University, Busan, Republic of Korea

<sup>3</sup>GenePath Dx, Shivajinagar, Pune – 411004, Maharashtra, India

\* Corresponding author: Dr. Parvathi A, CSIR-National Institute of Oceanography Regional Centre, Dr. Salim Ali Road, Post Box No. 1913, Kochi-682 018, India  
Ph: 91-(0)484-2390814, Fax: 91-(0)484-2390618 Email: parvathi@nio.org  
Running title: Bacterial diversity in Cochin estuary, India

### **Abstract**

In the present study, we analysed variations in bacterial community structure along a salinity gradient in a tropical monsoonal estuary (Cochin estuary, CE), on the southwest coast of India, using Illumina next-generation sequencing (NGS). Water samples were collected from eight different locations thrice a year, to assess the variability in the bacterial community structure and to determine the physico-chemical factors influencing the bacterial diversity. Proteobacteria was the most dominant phyla in the estuary followed by Bacteroidetes, Cyanobacteria, Actinobacteria, and Firmicutes. Statistical analysis indicated significant variations in bacterial communities between freshwater, mesohaline and euryhaline regions, as well as between the monsoon (wet) and non-monsoon (dry) periods. The abundance of Betaproteobacteria was higher in the freshwater regions, while Alphaproteobacteria and Epsilonproteobacteria were more abundant in mesohaline and euryhaline regions of the estuary. Gammaproteobacteria was more abundant in regions with high nutrient concentrations. Various bacterial genera indicating the presence of fecal contamination and eutrophication were detected. Corrplot based on Pearson correlation analysis demonstrated the important physico-chemical variables (Temperature, salinity, DO, and inorganic nutrients) that influence the distribution of dominant phyla, class and genera. The observed spatio-temporal variations in bacterial community structure in the CE were governed by regional variations in anthropogenic inputs and seasonal variations in monsoonal rainfall and tidal influx.

**Key words:** Cochin estuary, bacterial diversity, Next-generation sequencing (NGS), Estuaries Proteobacteria, monsoonal estuaries.

## **Introduction**

Marine microbial communities are vital to global biogeochemical cycles of carbon, nitrogen, sulphur, and phosphorous. They are the engines of every ecosystem and constitute massive biomass, diversity and activity in the global oceans (Graham et al. 2016). Hence, it is extremely important to understand the microbial community structure to appreciate how ecosystems function and to recognize factors that control microbial communities. But even with modern tools (high-throughput amplicon sequencing, metagenomics and metatranscriptomics), it is cumbersome to determine microbial community structure and map its variations in space and time. Physicochemical factors directly affect microbial diversity and community composition and variations in microbial community structure affect ecosystem functioning. Extensive studies have been carried out to reveal the bacterial diversity and community structure in many marine and lacustrine ecosystems (Bowman et al. 2003; Acinas et al. 2004; Heijs et al. 2008; Bobrova et al. 2016; Jeffries et al. 2016). However, only a few studies have addressed bacterial diversity from monsoonal estuaries (Crump et al. 2004; Bernhard et al. 2005; Khandeparker et al. 2017; Eswaran and Khandeparker 2019). Very little is known about the complex factors influencing bacterial community composition or the effects these communities have on estuarine ecosystems (Dolan 2005; Teira et al. 2008).

Estuaries make up some of the most complex and dynamic aquatic ecosystems due to freshwater influence from rivers and tidal influence from the seas. Mostly, the terrigenous riverine inputs together with the tidal mixing processes characterize the estuarine environments. CE is a highly dynamic tropical microtidal monsoonal estuary (Shivaprasad et al. 2013). The biodiversity in monsoonal estuaries is strongly influenced by monsoonal rains and riverine influx in addition to the estuarine variabilities in physical, chemical and biological factors due to tidal influx (Qasim 2003). The average monsoonal rainfall in the CE is 2038 mm (CWC data, 2016). The river influx from six major rivers amounts to 20000 mm<sup>3</sup>/year and the annual precipitation varied between 630 mm to 916 mm (Revichandran et al. 2012). During the monsoon (wet period), the riverine influx brings freshwater which accounts for 60–70% of the total annual river discharge to the system. The domestic sewage and industrial effluents dumped into the estuary results in nutrient enrichment in the CE (Madhu et al. 2007). During dry period (non-monsoonal months), the tidal influx is more pronounced due to reduced freshwater influx and precipitation (Madhu et al. 2007; Srinivas et al. 2003). Due to the variation in the monsoonal rains, tidal influx, riverine inputs, and the associated pollutants, the water quality of the ecosystem and the associated bacterial community diversity is affected (Vajravelu et al., 2018).

The bacterial community diversity in the CE as well as the impact of physicochemical parameters on the distribution of these communities concerning monsoonal rains have not been

studied so far. It is important to study the spatial and seasonal pattern in bacterial community structure as it reflects the selection mechanisms exerted by the dynamic environment on bacterial groups with specific functions and properties. Though microbial biogeography is addressed in many environments in recent years, principles that govern microbial distribution remain poorly understood (Thompson et al. 2017; Nemergut et al. 2011). Furthermore, metagenomic analysis of bacterial communities from estuarine environments found that salinity is the most important factor influencing bacterial composition in estuarine environments compared to other physico-chemical factors (Crump et al. 2004; Dong et al. 2004; Wu et al. 2019; Herfort et al. 2017). We hypothesised that the distribution of different bacterial groups in the CE could be a function of spatial gradients and seasonal variabilities in salinity and monsoonal rains.

## **Materials and Methods**

### ***Station description and sampling details***

The CE is a complex shallow estuary with an average depth of 4 m (**Fig. 1**). Six major rivers, the Pamba, Achancovil, Manimala, Meenachil, Periyar, and Muvattupuzha along with their tributaries and several canals bring large volumes of fresh water into the CE. The saline water from the neighbouring Arabian Sea enters the CE through the two inlets—one at Cochin and the other at Azhikode (**Fig. 1**). During the peak southwest monsoon (June–September), the rivers transport an enormous amount of freshwater into the CE, which transforms it almost entirely into a freshwater lake except near the two inlet regions. While in the dry period (non-monsoonal months, October-May), the riverine influxes gradually decrease, allowing salinity to build up in the estuary (Qasim 2003; Jyothibabu et al. 2006).

The water samples were collected from 8 distinct stations along the estuary during three months (August, November, and February in 2015-2016) (**Fig. 1**). The 8 stations were distinct with respect to inputs and outputs from the river and tides. Station 1 was located far upstream of the Periyar river, while Station 2 runs through the Industrial Belt of the city. Station 3 is the region where the Periyar river empties into the estuary, and Station 4 or Kochi inlet is where the estuary meets the Arabian Sea. Stations 5, 6, and 7 are located further downstream of Kochi, which receives a lot of sewage wastes from the urban population, similar to S3. Station 8 is situated beyond the Thanneermukkam tidal saltwater barrage near rice paddy plantations.

Water (5 L) samples were collected from the surface from each station in sterilized 1 L glass bottles, immediately placed on ice, and shielded from sunlight. At each station, the salinity, water temperature, and pH were recorded. The subsamples (triplicate) were collected to reduce the sampling

variability at each station during the study period. Upon returning to the lab, the samples were filtered through 0.2 µm filters (0.47 mm diameter, Millipore USA) using a sterilized vacuum filtration apparatus. Once filtration was complete, the filters were stored at -80°C until further processing.

### ***Environmental parameters***

Temperature and salinity were measured using a Conductivity Temperature Density profiler (CTD, SBE, Seabird 19). The inorganic nutrients (nitrate, nitrite, phosphate, ammonia, and silicate) were estimated spectrophotometrically (Shimadzu UV 1800) using standard protocols (Grasshoff 1983). Dissolved oxygen (DO) content was determined following Winkler's titration method (Grasshoff 1983).

### ***Enumeration of Total Plate count (TPC)***

Total plate count was employed to enumerate the viable bacteria in the water sample. Briefly, the water samples were serially diluted using 0.85% physiological saline and plated on to nutrient agar (NA). The plates were then incubated under room temperature for 24 hours. The colonies that developed on NA plates were enumerated and TPC was expressed as a number of colony forming units (cfu) /ml for a water sample.

### ***Extraction of DNA from water samples***

DNA was extracted from bacteria concentrated on 0.2 µm filters (Millipore, USA). Under aseptic conditions, the frozen filters were thawed, cut into small pieces using sterilized scissors. The total DNA was extracted using the Power-Soil DNA isolation kit following the manufacturer's instructions (Qiagen, USA) (Cao et al. 2013). The DNA (triplicate) from each sampling station was pooled and sent to GenePath Dx (Pune, India) for library construction and next-generation sequencing.

### ***Amplicon library construction***

The DNA samples were first quantified using a Broad Range Qubit System (Life Technologies, CA, USA), and the average concentration was 10 ng/µl. Library construction involved two PCR reactions. The first reaction ligated two 20 base pair (bp) proprietary tags on either end of the targeted V3-V4 region using 16S Illumina primers F: 5'- CCTACGGGNGGCWGCAG-3' and R: 5'- GACTACHVGGGTATCTAATCC-3' (Jasna et al. 2020, Parvathi et al. 2019) with the initial amount of 20 ng of DNA. Cycle conditions included an initial denaturation at 95°C for 3 min, followed by 40 cycles of 95 °C for 15 sec, 60 °C for 20 sec, 72 °C for 20 sec, followed by a final extension at 72 °C for 10 min (Klindworth et al. 2013). The amplified products of 400-600 bp were visualized by gel electrophoresis, diluted 1:10 using 10 mM Tris-HCl (pH 8.0), and used as templates for the second

PCR. This indexing PCR was completed using QuantiTect MultiPlex PCR kit (Qiagen, Germany). Both forward and reverse indexing primers contained a 100 bp tag, including the adapter and unique barcode sequences, and were used at 200 nM. Cycling conditions included, initial denaturation of 95 °C for 15 min followed by 30 cycles of denaturation at 95 °C for 30 sec, annealing at 63 °C for 45 sec, and extension at 72 °C for 90 sec. The amplified products were quantified using a Qubit Broad Range system (Life Technologies, CA, USA) (Parvathi et al. 2019; Ramanan et al. 2016).

### ***Illumina sequencing***

PCR products were molar-normalized, pooled into a single tube and purified to a minimum of 300 bp using PureLink PCR Purification kit (Invitrogen, CA, USA). The purified, pooled sample was then diluted to 4 nM final concentration using Resuspension Buffer (RSB – Illumina, CA, USA). The sample was denatured for 5 minutes and neutralized using 0.2 N NaOH and HT1 Buffer, respectively (Illumina, CA, USA). It was then pooled with other libraries prepared for NGS in a ratio dependent on amplicon size/total panel size, desired sequencing depth, and number of samples pooled in each sub-library. Pooled libraries were further diluted down to a final 15 pM and spiked with 5% phiX (Illumina, CA, USA) as a control and diversity enhancer. Samples were then loaded into an Illumina MiSeq v3 cartridge (Illumina, CA, USA) and run in 2\*300 mode on an Illumina MiSeq next generation sequencer (Illumina, CA, USA) (Parvathi et al. 2019; Ramanan et al. 2016).

### ***Initial processing of sequence reads***

The bcl2fastq Conversion Software embedded in the MiSeq was used for demultiplexing. The quality reads (>Q30) were filtered out using the automated FASTQ Tool Kit application on Illumina BaseSpace Labs website for downstream analysis. Within the application, TagCleaner software was used to remove the adapter sequences.

### ***Data analysis***

The QIIME pre-processing application (Version 1.0.0, Illumina BaseSpace) was used for the analysis of raw sequence data (Caporaso et al. 2010). Pre-processing included demultiplexing, quality filtering, OTU picking, and taxonomic assignment using Greengenes. QIIME outputs were used to create a BIOM file. Before succeeding analysis, OTUs abundance was normalized. QIIME software (version 1.7.0) was used to analyse alpha diversity and richness). The data were normalized and transformed by the Bray-Curtis method. PERMANOVA was performed to understand the significant variations in species abundance. A correlation matrix or correlogram was generated for analysing the relationship between environmental factors and dominant phyla, class and genus using the corrplot R package, version 0.84 (Wei and Simko 2017). Canonical correspondence analysis (CCA), a

multivariate method was used to elucidate the relationships between biological and environmental variables (CANOCO 4.5) (Lepš and Šmilauer 2003). A Monte Carlo test was used to determine the significance of each axis and to evaluate the influence of the environmental variables upon the overall distribution of bacterial species and their distribution at each sites and sampling seasons. (Salles et al. 2004; Sapp et al. 2007). Similarity percentage breakdown (SIMPER) was used to calculate the partition of the average Bray-Curtis dissimilarity between different clusters into components from different genera (Clarke 1993). This allowed the identification of genera that are the most important in creating observed patterns in similarity.

### ***Nucleotide sequence accession number***

Paired end Illumina sequence data from this study was submitted to the NCBI Sequence Read Archive (SRA) under bio project number PRJNA595444.

## **Results**

### ***Environmental parameters***

Based on the climatology of the study area, seasons have traditionally been classified into monsoon (June to September), post-monsoon (October to January), and pre-monsoon (February to May) (Menon et al. 2000). Maximum rainfall, accounting for 60–65 % of the total annual rainfall in the study area, was received from June to September (wet period, with an average rainfall of 528.75 mm), whereas the dry period, sub-classified into the Dry period I and II, received an average rainfall of 151.75 mm and 133 mm, respectively (**Fig. S1**). Salinity was the most fluctuating variable in the CE ranging from 0-28. The entire estuary was freshwater dominated during the wet period, except at the inlet station, S4, which sustained a salinity of 16. With the retreat of the southwest monsoon, the riverine influx reduced by 45% resulting in salinity stratification in the estuary with euryhaline salinity at the inlet (**Table 1**), freshwater conditions in stations S1, S2, S7 and S8 and mesohaline salinities in stations S3, S5, and S6 (**Table 1**). The major inorganic nutrients were high during the sampling period in the CE (**Table 1**).

### ***Total Plate Count, Taxonomic richness and $\alpha$ -diversity of the prokaryotic community***

Total plate count (TPC) of bacteria ranged from 0.22 to  $5.98 \times 10^5$  cfu/ml (**Fig S2**). The bacterial abundance was higher in the Dry period II and especially in the euryhaline region of the estuary. The bacterial community diversity was highest in the dry period I compared to other periods (**Table 2**). The overall species coverage of the samples was highest ( $\geq 68.60\%$ ) for dry period I, followed by wet period ( $\geq 66.68\%$ ) and the dry period II ( $\geq 63.19\%$ ). The Good's average was high (0.99) indicating

that most of the bacterial diversity was attained by sequencing. However, Shannon diversity, Chao richness and Simpson diversity indices, which represent abundance and evenness of the species distribution, varied slightly with sampling period and with stations (**Table 2**). There was a significant difference in  $\alpha$ -diversity between the dry and wet seasons at all stations (Kruskal–Wallis,  $p < 0.05$ ) and bacterial diversity significantly differed both spatially and temporally (Kruskal–Wallis,  $p < 0.05$ ).

### ***Seasonal and spatial variations in bacterial diversity at phylum and class level***

The relative abundances of different phyla in the eight locations from three different sampling periods (24 samples) are shown in **Fig. 2**. Total bacterial diversity was distributed among 32 different phyla, with Proteobacteria, Bacteroidetes, Firmicutes, Actinobacteria, Cyanobacteria, and Verrucomicrobiae accounting for more than  $> 95\%$  of the total OTUs in all the samples (**Fig. 2**). Phylum Proteobacteria was dominated by Alphaproteobacteria, Gammaproteobacteria, and Betaproteobacteria, with Betaproteobacteria being more abundant in the freshwater regions and Alphaproteobacteria in the meso- and euryhaline regions of the estuary. Betaproteobacteria was the most dominant class throughout the freshwater-dominated estuary during the wet period, in all stations except S4 (**Fig. S3**).

### ***Bacterial diversity at generic level***

A total of 975 genera contributing to 60 to 75 % of total generic diversity was considered and represented in **Fig. 3**. The most dominant genus was *Sanguibacter*, *Saccharopolyspora*, *Prochlorococcus*, *Arcobacter*, and *Ruegeria*, with significant spatial variations in abundance. The dominant genus was distinctly different in the freshwater, mesohaline and euryhaline regions of the estuary. Most dominant genus in the freshwater regions were *Sanguibacter*, *Saccharopolyspora*, *Demequina*, *Chthoniobacter*, *Bifidobacterium*, *Paucibacter*, *Flavobacterium*, *Limnohabitans*, *Chitinophaga*, *Acinetobacter*, *Oxalobacter*, *Prochlorococcus*, etc. In Mesohaline regions, dominant genera included *Microcystis*, *Prochlorococcus*, *Agromyces*, *Saccharopolyspora*, *Actinocatenispora*, *Peptoniphilus*, *Acidiphilium*, *Polaribacter*, *Aquimarina*, *Arcobacter*, *Gramella*, *Leucobacter*, *Flavobacterium*, *Ruegeria*, *Calothrix* etc. *Actinocatenispora*, *Agromyces*, *Prochlorococcus*, *Microcystis*, *Acidiphilium*, *Ruegeria*, *Saccharopolyspora*, *Vibrio*, *Peptoniphilus*, *Gramella*, *Aquimarina*, *Leucobacter*, *Acidocella*, *Tenacibaculum*, *Marivita*, *Nisaea*, *Arcobacter*, were most dominant in the euryhaline regions of the estuary. Many bacterial genera in the mesohaline regions were also present in the euryhaline regions. However, their percentage abundance varied in these two regions of the estuary. Pollution indicators such as *Bacteroides*, *Microcystis*, *Agromyces*, *Vibrio*, *Clostridium*, *Prevotella*, *Enterobacter*, *Ruminococcus*, *Lachnospira*, *Pseudomonas* were detected.

## *Statistical Analysis*

NMDS analysis revealed two clusters of bacterial communities during wet period (**Fig. 4**). During wet period, the entire estuary considered like a freshwater lake and hence formed two clusters, S4 in one cluster and all other stations in another cluster. During dry months, the bacterial communities were clustered into three, euryhaline stations (S4, salinity = >25), mesohaline stations (S3, S5, and S6, salinity=5-20) and freshwater stations comprising stations S1, S2 and S8 (salinity =<5). Based on SIMPER analysis, the average similarity within the cluster was 73 % and the degree of dissimilarity between freshwater and mesohaline clusters was 40 %. Dry period 1 yielded three clusters with an average similarity of 75% in the freshwater cluster. The average dissimilarity between freshwater and mesohaline stations was 32% and that between freshwater and euryhaline stations was 38%. The average dissimilarity between mesohaline and euryhaline stations was 26%. NMDS revealed two clusters in dry period II. The average dissimilarity between freshwater and mesohaline stations was 42% and that between mesohaline and euryhaline stations was 21% in dry period II.

CCA analysis of the phylum, class, and generic level diversity showed a distinct spatial and seasonal pattern in the distribution of bacterial communities (**Fig. S4**). Distribution of bacterial communities at each station was influenced by different environmental parameters, such as DO, silicate, and nitrate in the freshwater regions, high inorganic nutrients such as ammonia, phosphate, and nitrite in the mesohaline regions and salinity in the euryhaline station, S4. Correlogram based on Pearson correlation analysis revealed the influence of physico-chemical parameters on the distribution of dominant phyla, class, and genus (**Fig. 5**). Proteobacteria, Cyanobacteria and Alphaproteobacteria showed a positive correlation with nitrite, and ammonia. Betaproteobacteria showed a positive correlation with silicate, DO, nitrate and temperature (**Fig. 5**). Salinity and other physico-chemical factors played an important role in determining the distribution of Cyanobacteria, Gammaproteobacteria, Alphaproteobacteria and dominant bacterial genera in the estuary (**Fig. 5**).

The number of shared and unique OTUs at the genus level in dry and wet periods is indicated in the Venn diagram (**Fig. S5**). Comparative analysis showed that ~1621 OTUs were shared during all the seasons, 172 OTUs were unique to wet period, and 375 OTUs to dry period I and 87 OTUs to dry period II. PERMANOVA analysis demonstrated that bacterial community structure significantly varied temporally during wet and dry seasons (Pseudo F = 3.342,  $p = 0.002$ ) and spatially (Pseudo F = 1.615,  $p = 0.016$ ).



## Discussion

Estuaries, being dynamic mixing zones of the ocean and freshwater masses, are characterized by steep spatial and temporal gradients of physical, chemical, and biological parameters. Hence, it is essential to fathom the impact of these gradients on the local bacterial community, their metabolism and the water quality in an estuarine system. Salinity has been demonstrated as an important environmental factor structuring bacterial communities in coastal ecosystems (Ortega-Retuerta et al. 2013; Liu et al. 2015; Herlemann et al. 2016). Though CE is comprised of similar aquatic microbial phyla found within other aquatic environments (Eswaran and Khandeparker 2019; Meziti et al. 2016; Savio et al. 2015), it hosted variations in the relative abundance of bacterial communities with changes in estuarine hydrography and pollutants. The estuarine bacterial genera fell into three distinct categories, Euryhaline/marine (Salinity > 20), mesohaline/brackish (Salinity = 5-20), and freshwater (Salinity = <5) (See Station details and description for seasonal hydrography in the CE) (**Fig. 4**). Previous studies in the CE have demonstrated the existence of three distinct zones based on salinity variations during dry months and two zones during wet monsoonal months and have indicated unique biological responses to these gradients (Parvathi et al. 2015; Jasna et al. 2017). In addition to the variations imparted by salinity influx, there are regional variations in the input of industrial wastes, agricultural wastes, and sewage inputs, especially from many non-point sources. Large inputs of fresh organic matter into the estuary from riverine inputs and other non-point sources impact heterotrophic production (Jasna et al. 2017) and bacterial diversity. We detected fluctuations in bacterial richness within the three different salinity regimes of the estuary (**Table 2**). This indicates that factors other than salinity influenced bacterial richness. Freshwater regions of the estuary detected the highest and lowest richness during dry period. We assume that variations in the lability of organic matter in these regions might be responsible for the proliferation of several adapted bacterial taxa (Bunse et al. 2016). The results of this study corroborated with recent studies on distinct bacterial communities in estuarine environments of Delaware Bay, Chesapeake Bay, Columbia River estuary, and Baltic Sea (Herfort et al. 2017; Herlemann et al. 2011; Hugerth et al. 2015).

The dominance of Proteobacteria in the present study was in concurrence with reports from other tropical estuaries (Bouvier and del Giorgio 2002; Ghosh and Bhadury 2019; Ortmann and Santos 2016; Eswaran and Khandeparker 2019) and also from coastal waters of India (Sachithanandam et al. 2020; Parvathi et al. 2019). The high abundance of Betaproteobacteria and Alphaproteobacteria in our study corroborated with previous studies from other estuarine regions (Sekiguchi et al. 2002). While, Betaproteobacteria was found in high proportions in the freshwater regions of the estuary, Alphaproteobacteria, Gammaproteobacteria and Epsilonbacteria were more dominant in the

mesohaline and euryhaline regions of the estuary during dry seasons. Betaproteobacteria was present in high proportions at salinities below 4 and Alphaproteobacteria at salinities above 13. This shows that salinity transitions play a significant role in the abundance and distribution of Beta- and Alphaproteobacteria. However, the mechanisms causing changes in bacterial community composition at different salinities are currently unclear. Bacteria have remarkable versatility to adapt to different trophic conditions and hence, environmental variabilities directly reflect upon the distribution and abundance of bacterial communities. Few studies in the CE have linked salinity to differences in the key metabolic capabilities of bacteria (Thottathil et al. 2008 a).

Dominance of Proteobacteria, Cyanobacteria, Alphaproteobacteria, and Epsilonproteobacteria was high in regions with high concentrations of ammonia and nitrite. Mesohaline regions were characterized by high concentrations of ammonia, nitrite, and phosphate. The mesohaline regions of the estuary received a high amount of agriculture wastes, industrial effluents, and sewage from the urban population through many nonpoint sources. This region has the highest amount of ammonia and nitrite indicating increased sewage input and agricultural runoff. These regions lie between the inlet and the freshwater region/Vembanad Lake, which makes these regions less dynamic with low flushing rates during the dry seasons with no rainfall and less riverine influx (Jasna et al. 2017). Dissolved organic carbon is also more ( $340\pm 108\ \mu\text{M}$  to  $193\pm 102\ \mu\text{M}$ ) in the mesohaline regions compared to freshwater regions ( $<200\ \mu\text{M}$ ) of the estuary (Gupta et al. 2009). Moreover, due to the closure of Thannermukkam bund during dry period II, the riverine influx from the Vembanad Lake is blocked to prevent salinity intrusion to the paddy fields in the southern regions (**Fig 1**). This has reduced the flushing out of organic pollutants from the estuary, especially from these less dynamic mesohaline regions. The mesohaline regions support high bacterial respiration (Jasna et al. 2017), indicating that these regions have a heavy nutrient load which in turn supports the growth and dominance of Gammaproteobacteria, Alphaproteobacteria, and Epsilonproteobacteria.

Saline water allows the dissolution of dissolved inorganic phosphorus (DIP) from iron-bound fractions in sediment, leading to a subsequent increase in phosphate observed in surface waters (Slomp 2011). On the contrary during wet season, river discharge brings DIP into estuaries from anthropogenic as well as natural sources (Slomp 2011). Phosphate is a major constituent of nucleic acid and lipids, hence is an important nutrient required for the growth of microorganisms. Gammaproteobacteria were also abundant in the river mouth station, S1 during wet season, indicating that this group also takes advantage of allochthonous material loadings and nutrient-enriched conditions. In addition to these, variations in other biotic factors, such as high phytoplankton (Madhu et al. 2007), grazing (Sooria et al. 2016), and viral lysis (Jasna et al. 2017) can also shape bacterial community composition in the CE.

For eg., Diatom have been the primary determinant for *Vibrio* population in the coastal water of the Arabian sea along the southwest coast of India (Asplund et al. 2011). The growth of autochthonous bacteria, for instance, *Vibrio* might be supported by diatom-dominated phytoplankton as well as zooplankton communities in the CE. The results of our study point to the potential that *Vibrio* strains are sustained as free-living populations in the CE. Hence the observed variations in bacterial community composition patterns may be a result of a multitude of factors influenced by trophic conditions, anthropogenic inputs, riverine inputs, and tidal influx.

The presence of some genera indicated the extent of pollution in this estuarine environment. Some strains of genus *Sanguibacter*, *Bifidobacterium*, *Oxalobacter* are known to colonize the animal gastrointestinal tract and in rumens of animals such as cattle. They are also isolated from marine and freshwater environments (Garrity et al. 2005). The most dominant genus in mesohaline regions was *Microcystis* which include members that can produce neurotoxins and hepatotoxins, such as microcystin and cyanopeptolin. The presence of these pathogens in the CE indicates the presence of sewage or fecal contamination from other animal sources. *Bacteroides* are an important indicator group which are exclusive to warm-blooded animals and constitutes a larger portion of pollution indicator bacteria in CE, especially during dry period. Human and animal associated *Bacteroides* markers are been extensively used for faecal identification studies (Harwood et al. 2014). In the present study, *Bacteroides* showed a significant positive correlation with nitrate and ammonia and did not show any correlation with salinity. Anthropogenic nutrient over-enrichment (eutrophication), and other factors including expansion of intensive agriculture, rapid industrialization, and urbanization, enhance the occurrence of microcystin-producing HABs in most regions and thus poses deleterious effects on human health (Garrity et al. 2020). Bacteria involved in faecal contamination such as *Bacteroides*, *Clostridium*, *Faecalibacterium*, *Enterococcus*, *Pseudomonas*, *Vibrio*, *Prevotella*, *Enterobacter*, *Klebsiella*, *Campylobacter* are also detected in CE (Boehm and Sassoubre 2014).

Several genera in the CE could be considered as indicators of eutrophication, such as those involved in ammonia oxidation, nitrite reduction, N<sub>2</sub>O reduction. Genus such as *Paracoccus*, *Comamonas*, *Nitrosomonas*, and *Nitrobacter* are involved in the nitrogen cycle and were more abundant in the mesohaline regions of the estuary (Wang et al. 2014; Kim et al. 2015). Dissolve inorganic nitrogen is high in the CE with a higher nitrogen fixation rate (Bhavaya et al. 2016). These regions receive 7-11 times high organic matter as lateral input as compared to the inputs through rivers (Thottathil et al. 2008b; Gupta et al. 2009). The input of industrial and domestic waste discharges as a result of large human settlement and industrial growth and agriculture has resulted in excess nutrients in the system. The high abundance of *Prochlorococcus marinus* and *Synechococcus* can be

considered as an indicator of trophic status of the CE (Rajaneesh et al. 2015). The trophic index scores (TRIX units) showed that CE is highly eutrophic (Martin et al. 2012, Hershey et al. 2019). Hence, it is important to introduce proper management measures to impede the nutrient loading from various non-point to protect the health of this estuary.

## **Conclusion**

Our study demonstrated that the seasonal and spatial variations in bacterial community structure are largely influenced by salinity and inorganic nutrients like nitrite, nitrate, ammonia, phosphate, and silicate in the CE. Spatial variations in bacterial community structure were also influenced by regional variations in anthropogenic inputs to a large extent. Regional variations in anthropogenic inputs, especially in the mesohaline regions, imparted by restricted flow in the CE, had a greater impact in shaping the bacterial community structure during dry periods. Our study demonstrates how bacterial community structure changes along an estuarine gradient during different seasons and the physico-chemical conditions that drive bacterial community shifts. Proteobacteria was the most dominant bacterial phylum, mainly composed of the classes Alphaproteobacteria, Betaproteobacteria, and Gammaproteobacteria. Various genera indicative of eutrophication, fecal contamination, and sewage pollution were identified in this study. Our study suggests that monitoring the presence of important bacterial groups could serve as an appropriate indicator of ecosystem health and pollution. This estuarine system experiences elevated stress from human activity and increased knowledge of factors that shape its microbiology is crucial.

## **Acknowledgements**

The authors are grateful to the Director, NIO, Goa and all our colleagues in CSIR-NIO (RC), Kochi for their support and advice. JV is grateful to Council of Scientific and Industrial Research (CSIR), New Delhi, for the senior research fellowship grant. MC is grateful to Fulbright Nehru Fellowship (USIEF). We are thankful to Dr Abdul Jaleel, Scientist, CSIR-NIO for guidance in statistical analysis. We are grateful to Mr. Vishal CR, Research Scholar, CSIR-NIO for helping us the CANOCO analysis. We are grateful to Dr. Deepak George Pazhayamadom, Museum and Art Gallery Northern Territory, Darwin, Australia, for corrplot analysis using R. This is NIO contribution number xxxx.

## **Funding**

JV is grateful to Council of Scientific and Industrial Research (CSIR), New Delhi, for the senior research fellowship grant. MC is grateful to the Fulbright Nehru Fellowship (USIEF).

#### **Conflicts of interest/Competing interests**

Authors declare no conflicting or competing interests

#### **Ethics approval**

No need of ethics approval, since this work does not include research on identifiable human material or data.

#### **Consent to participate**

All the authors agree to participate in this manuscript

#### **Consent for publication**

The manuscript has written consent form all the authors and from the institution.

#### **Availability of data and materials**

The data that support the findings of this study are available from the corresponding author upon reasonable request.

#### **Credit authorship contribution statement**

AP conceptualized the work, made funding arrangements data analysis and prepared the manuscript. MC conceptualisation, data collection and sample analysis, JV analysed samples, performed statistical analysis and prepared the manuscript. NP, NG carried out the Illumina Miseq sequencing.

#### **References**

- Acinas SG, Klepac-Ceraj V, Hunt DE, Pharino C, Ceraj I, Distel DL, Polz MF. (2004). Fine-scale phylogenetic architecture of a complex bacterial community. *Nature*, 430:551. <https://doi.org/10.1038/nature02649>
- Asplund, M. E., Rehnstam-Holm, A. S., Atnur, V., Raghunath, P., Saravanan, V., Härnström, K., ... & Godhe, A. (2011). Water column dynamics of *Vibrio* in relation to phytoplankton community composition and environmental conditions in a tropical coastal area. *Environ Microbiol.* 13(10), 2738-2751. <https://doi.org/10.1111/j.1462-2920.2011.02545.x>
- Bernhard AE, Donn T, Giblin AE, Stahl DA. (2005). Loss of diversity of ammonia-oxidizing bacteria correlates with increasing salinity in an estuary system. *Environ. Microbiol.* 7:1289-1297. <https://doi.org/10.1111/j.1462-2920.2005.00808.x>
- Bhavya PS, Kumar S, Gupta GVM, Sudheesh V, Sudharma KV, Varrier DS, ... & Saravanane N. (2016). Nitrogen uptake dynamics in a tropical eutrophic estuary (Cochin, India) and adjacent coastal waters. *Estuaries and Coasts*, 39(1), 54-67. <https://doi.org/10.1007/s12237-015-9982-y>
- Bobrova O, Kristoffersen JB, Oulas A, Ivanytsia V. (2016) Metagenomic 16s rRNA investigation of microbial communities in the Black Sea estuaries in South-West of Ukraine. *Acta Biochim Pol.*63(2):315-319. DOI: [https://doi.org/10.18388/abp.2015\\_1145](https://doi.org/10.18388/abp.2015_1145)
- Boehm AB, Sassoubre LM. (2014). Enterococci as indicators of environmental fecal contamination. *Enterococci: from commensals to leading causes of drug resistant infection [Internet]*.
- Bouvier TC, del Giorgio PA. (2002). Compositional changes in free-living bacterial communities along a salinity gradient in two temperate estuaries. *Limnol. Oceanol.*, 47:453-470. <https://doi.org/10.4319/lo.2002.47.2.0453>
- Bowman JP, McCammon SA, Gibson JA, Robertson L, Nichols PD. (2003). Prokaryotic metabolic activity and community structure in Antarctic continental shelf sediments. *Appl. Environ. Microbiol.* 69:2448–2462. DOI: [10.1128/AEM.69.5.2448-2462.2003](https://doi.org/10.1128/AEM.69.5.2448-2462.2003)
- Bunse C, Bertos-Fortis M, Sassenhagen I, Sildever S, Sjöqvist C, Godhe A, ... & Legrand, C. (2016). Spatio-temporal interdependence of bacteria and phytoplankton during a Baltic Sea spring bloom. *Front. Microbiol.* 7, 517. <https://doi.org/10.3389/fmicb.2016.00517>
- Cao Y, Van De Werfhorst LC, Dubinsky EA, Badgley BD, Sadowsky MJ, Andersen GL, Griffith JF, Holden PA. (2013). Evaluation of molecular community analysis methods for discerning fecal sources and human waste. *Water Res.* 47: 6862-6872. <https://doi.org/10.1016/j.watres.2013.02.061>
- Clarke KR. (1993). Non-parametric multivariate analyses of changes in community structure. *Australian Journal Ecol.*, 18(1), 117-143. <https://doi.org/10.1111/j.1442-9993.1993.tb00438.x>
- Crump BC, Hopkinson CS, Sogin ML, Hobbie JE. (2004). Microbial biogeography along an estuarine salinity gradient: combined influences of bacterial growth and residence time. *Appl. Environ. Microbiol.*, 70:1494-1505. DOI: [10.1128/AEM.70.3.1494-1505.2004](https://doi.org/10.1128/AEM.70.3.1494-1505.2004)

- Dolan JR. (2005). An introduction to the biogeography of aquatic microbes. *Aquatic Microb. Ecol.*, 41:39-48. [doi:10.3354/ame041039](https://doi.org/10.3354/ame041039)
- Dong L, Su J, Wong LA, Cao Z, Chen JC. (2004). Seasonal variation and dynamics of the Pearl River plume. *Cont. Shelf. Res.* 24:1761–1777. <https://doi.org/10.1016/j.csr.2004.06.006>
- Eiler A, Carlos Gonzalez-Rey, Sophie Allen, Stefan Bertilsson, Growth response of *Vibrio cholerae* and other *Vibrio* spp. to cyanobacterial dissolved organic matter and temperature in brackish water, *FEMS Microbiol Ecol*, 60: 411–418, <https://doi.org/10.1111/j.1574-6941.2007.00303.x>
- Eswaran R, Khandeparker L. (2019). Seasonal variation in  $\beta$ -glucosidase-producing culturable bacterial diversity in a monsoon-influenced tropical estuary. *Environ Monitor Assess*, 191(11), 1-11. <https://doi.org/10.1007/s10661-019-7818-0>
- Garrity GM, Bell JA, Lilburn T. (2005). The revised road map to the Manual. In *Bergey's Manual of Systematic Bacteriology*, 2nd edn, vol. 2A, pp. 159–187. Edited by D. J. Brenner, N. R. Krieg & J. T. Staley. New York: Springer
- Garrity, George Jitendra K. Malik, Vijay K. Bharti, Anu Rahal, Dinesh Kumar, Ramesh C. Gupta, Chapter 31 - Cyanobacterial (blue-green algae) toxins, Editor(s): Ramesh C. Gupta, *Handbook of Toxicology of Chemical Warfare Agents (Third Edition)*, Academic Press, 2020, Pages 467-478, ISBN 9780128190906
- Ghosh A, Bhadury P. (2019). Exploring biogeographic patterns of bacterioplankton communities across global estuaries. *Microbiol. Open.* 8(5):e00741. <https://doi.org/10.1002/mbo3.741>
- Grasshoff K, Ehrhardt MKK. (1983). In K. Grasshoff, K. Kremling, & M.Ehrhardt (Eds.), *Methods of seawater analysis* (pp. 89–224). Weinheim: Verlag Chemie.
- Gupta GVM, Thottathil SD, Balachandran KK. et al. (2009) CO<sub>2</sub> Supersaturation and Net Heterotrophy in a Tropical Estuary (Cochin, India): Influence of Anthropogenic Effect. *Ecosystems* 12:1145–1157. <https://doi.org/10.1007/s10021-009-9280-2>
- Harwood VJ, Staley C, Badgley BD, Borges K, Korajkic A (2014) Microbial source tracking markers for detection of fecal contamination in environmental waters: relationships between pathogens and human health outcomes. *FEMS Microbiol Rev* 38:1–40. <https://doi.org/10.1111/1574-6976.12031>
- Heijs SK, Laverman AM, Forney LJ, Hardoim PR, Van Elsas JD. (2008). Comparison of deep-sea sediment microbial communities in the Eastern Mediterranean. *FEMS Microbiol Ecol*, 64:362-377. <https://doi.org/10.1111/j.1574-6941.2008.00463.x>
- Herfort L, Crump BC, Fortunato CS, McCue LA, Campbell V, Simon HM, ... & Zuber P. (2017). Factors affecting the bacterial community composition and heterotrophic production of Columbia River estuarine turbidity maxima. *Microbiol. Open*, 6(6), e00522. <https://doi.org/10.1002/mbo3.522>
- Herlemann DP, Lundin D, Andersson AF, Labrenz M, Jürgens K. (2016). Phylogenetic signals of salinity and season in bacterial community composition across the salinity gradient of the Baltic Sea. *Front. Microbiol.* 7:1883. <https://doi.org/10.3389/fmicb.2016.01883>

- Herlemann DP, Labrenz M, Jürgens K, Bertilsson S, Waniek JJ, Andersson AF. (2011). Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME J.*, 5(10), 1571-1579. <https://doi.org/10.1111/1462-2920.12886>
- Hugerth LW, Larsson J, Alneberg J, Lindh MV, Legrand C, Pinhassi J, & Andersson AF. (2015). Metagenome-assembled genomes uncover a global brackish microbiome. *Genome boil.*, 16(1), 1-18. <https://doi.org/10.1186/s13059-015-0834-7>
- Jasna V, Parvathi A, Pradeep Ram AS, Balachandran KK, Madhu NV, Nair M, ... & Sime- Ngando T. (2017). Viral-induced mortality of prokaryotes in a tropical monsoonal estuary. *Front Microbiol.*, 8:895 <https://doi.org/10.3389/fmicb.2017.00895>
- Jeffries TC, Schmitz Fontes ML, Harrison DP, Van-Dongen-Vogels V, Eyre BD, Ralph PJ, & Seymour JR. (2016). Bacterioplankton dynamics within a large anthropogenically impacted urban estuary. *Front Microbiol.*, 6: 1438. <https://doi.org/10.3389/fmicb.2015.01438>
- Jyothibabu R, Madhu NV, Jayalakshmi KV, Balachandran KK, Shiyas CA, Martin GD, Nair KK. (2006). Impact of freshwater influx on microzooplankton mediated food web in a tropical estuary (Cochin backwaters–India). *Estuarine, Coastal Shelf Sci.*, 69:505-518. <https://doi.org/10.1016/j.ecss.2006.05.013>
- Khandeparker L, Eswaran R, Gardade L, Kuchi N, Mapari K, Naik SD, & Anil AC. (2017). Elucidation of the tidal influence on bacterial populations in a monsoon influenced estuary through simultaneous observations. *Environ. Monit. Assess.*, 189(1), 41. <https://doi.org/10.1007/s10661-016-5687-3>
- Kim H, Kaown D, Mayer B, Lee JY, Hyun Y, & Lee KK. (2015). Identifying the sources of nitrate contamination of groundwater in an agricultural area (Haeon basin, Korea) using isotope and microbial community analyses. *Sci. Total Environ.*, 533, 566-575. <https://doi.org/10.1016/j.scitotenv.2015.06.080>
- Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, Glöckner FO. (2013). Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res.*, 41e1-e1. <https://doi.org/10.1093/nar/gks808>
- Lepš, J., & Šmilauer, P. (2003). Multivariate analysis of ecological data using *CANOCO*. Cambridge university press.
- Liu J, Fu B, Yang H, Zhao M, He B, & Zhang XH. (2015). Phylogenetic shifts of bacterioplankton community composition along the Pearl estuary: The potential impact of hypoxia and nutrients. *Front. Microbiol.*, 6 article 64. <https://doi.org/10.3389/fmicb.2015.00064>
- Madhu NV, Jyothibabu R, Balachandran KK, Honey UK, Martin GD, Vijay JG, Shiyas CA, Gupta GV, Achuthankutty CT. (2007) Monsoonal impact on planktonic standing stock and abundance in a tropical estuary (Cochin backwaters–India). *Estuarine, Coastal Shelf Sci.* 1;73(1-2):54-64. <https://doi.org/10.1016/j.ecss.2006.12.009>
- Martin GD, George R, Shaiju P, Muraleedharan KR, Nair SM, Chandramohanakumar N. (2012). Toxic metals enrichment in the surficial sediments of a eutrophic tropical estuary (Cochin



Backwaters, Southwest Coast of India). *The Scientific World Journal*, 2012. <https://doi.org/10.1100/2012/972839>.

Menon NN, Balchand AN, Menon NR. (2000). Hydrobiology of the Cochin backwater system—a review. *Hydrobiologia*, 430(1), 149-183. <https://doi.org/10.1023/A:1004033400255>

Meziti A, Tsementzi D, Ar. Kormas K, Karayanni H, & Konstantinidis KT. (2016). Anthropogenic effects on bacterial diversity and function along a river-to-estuary gradient in Northwest Greece revealed by metagenomics. *Environ Microbiol.*, 18(12), 4640-4652. <https://doi.org/10.1111/1462-2920.13303>

Nemergut DR, Costello EK, Hamady M, Lozupone C, Jiang L, Schmidt SK., ... & Knight R. (2011). Global patterns in the biogeography of bacterial taxa. *Environ Microbiol.*, 13:1, 135-144. <https://doi.org/10.1111/j.1462-2920.2010.02315.x>

Ortega-Retuerta E, Joux, F, Jeffrey WH, & Ghiglione JF. (2013). Spatial variability of particle-attached and free-living bacterial diversity in surface waters from the Mackenzie River to the Beaufort Sea (Canadian Arctic). *Biogeosci*, 10, 2747– 2759. <https://doi.org/10.5194/bg-10-2747-2013>

Ortmann AC, Santos TT. (2016). Spatial and temporal patterns in the Pelagibacteraceae across an estuarine gradient. *FEMS Microb. Ecol*, 92: fiw133. <https://doi.org/10.1093/femsec/fiw133>

Parvathi A, Jasna V, Aswathy VK, Nathan VK, Aparna S, Balachandran KK. (2019). Microbial diversity in a coastal environment with co-existing upwelling and mud-banks along the south west coast of India. *Molecul. Biol. Rep.* 46:3113-27 <https://doi.org/10.1007/s11033-019-04766-y>

Parvathi A, Jasna V, Jina S, Jayalakshmy KV, Lallu KR, Madhu NV, ... & Balachandran K. (2015). Effects of hydrography on the distribution of bacteria and virus in Cochin estuary, India. *Ecol. Res.*, 30(1), 85-92. <https://doi.org/10.1007/s11284-014-1214-6>.

Qasim SZ. (2003). *Indian estuaries*. Allied publishers.

Ramanan V, Kelkar K, Ranade S, Gangodkar P, Gogate N, Patil K, Ragte-Wathare T, Agarwal M, Phadke ND. (2016). The clinical utility of a custom-developed targeted next-generation sequencing assay for detection of mutations associated with Philadelphia-negative chronic myeloproliferative neoplasms: Two case examples with CALR exon 9 mutations. *Int. Journal*, 1:29.

Rajaneesh KM, Smita M, Anil AC, Sawant SS. (2015). *Synechococcus* as an indicator of trophic status in the Cochin backwaters, west coast of India. *Ecol Indicator*. 55: 118-130 <https://doi.org/10.1016/j.ecolind.2015.02.033>

Revichandran C, Srinivas K, Muraleedharan KR, Rafeeq M, Amaravayal S, Vijayakumar K, & Jayalakshmy KV (2012). Environmental set-up and tidal propagation in a tropical estuary with dual connection to the sea (SW Coast of India). *Environ Earth Sci*, 66:4, 1031-1042. <https://doi.org/10.1007/s12665-011-1309-0>

- Sachithanandam V, Saravanane N, Chandrasekar K, Karthick P, Lalitha, P, Elangovan SS, & Sudhakar M. (2020). Microbial diversity from the continental shelf regions of the Eastern Arabian Sea: A metagenomic approach. *Saudi journal boil. Scie*, 27(8), 2065-2075. <https://doi.org/10.1016/j.sjbs.2020.06.011>
- Salles JF, Van Veen JA, & Van Elsas JD. (2004). Multivariate analyses of Burkholderia species in soil: effect of crop and land use history. *Appl. Environ. Microbiol*, 70(7), 4012-4020. DOI: [10.1128/AEM.70.7.4012-4020.2004](https://doi.org/10.1128/AEM.70.7.4012-4020.2004)
- Sapp M, Schwaderer AS, Wiltshire KH, Hoppe HG, Gerdt G, & Wichels A. (2007). Species-specific bacterial communities in the phycosphere of microalgae?. *Microb. Ecol.*, 53(4), 683-699. DOI: [10.1007/s00248-006-9162-5](https://doi.org/10.1007/s00248-006-9162-5)
- Savio D, Sinclair L, Ijaz UZ, Parajka, J, Reischer GH, Stadler P, ... & Eiler A. (2015). Bacterial diversity along a 2600 km river continuum. *Environ. microbial.*, 17(12), 4994-5007. <https://doi.org/10.1111/1462-2920.12886>
- Sekiguchi H, Watanabe M, Nakahara T, Xu B, Uchiyama H. (2002). Succession of bacterial community structure along the Changjiang River determined by denaturing gradient gel electrophoresis and clone library analysis. *Appl. Environ. Microbiol.* 68: 5142–5150. DOI: [10.1128/AEM.68.10.5142-5150.2002](https://doi.org/10.1128/AEM.68.10.5142-5150.2002)
- Shivaprasad A, Vinita J, Revichandran C, Reny PD, Deepak MP, Muraleedharan KR, NaveenKumar KR. (2013). Seasonal stratification and property distributions in a tropical estuary (Cochin estuary, west coast, India). *Hydrol. Earth Syst. Sci.* 17: 187–199. <https://doi.org/10.5194/hess-17-187-2013>
- Slomp CP (2011). Phosphorus Cycling in the Estuarine and Coastal Zones: Sources, Sinks, and Transformations, Editor(s): Eric Wolanski, Donald McLusky, Treatise on Estuarine and Coastal Science, Academic Press, 2011, Pages 201-229.
- Sooria PM, Jyothibabu R, Anjusha A, Vineetha G, Vinita J, Lallu KR, et al. (2015). Plankton food web and its seasonal dynamics in a large monsoonal estuary (Cochin backwaters, India) - Significance of mesohaline region. *Environ. Monit. Assess.* 187, 1–22. doi: [10.1007/s10661-015-4656-6](https://doi.org/10.1007/s10661-015-4656-6).
- Srinivas K, Revichandran C, Maheswaran PA, Asharaf TT, Murukesh N. (2003) Propagation of tides in the Cochin estuarine system, southwest coast of India.
- Teira E, Gasol JM, Aranguren-Gassis M, Fernández A, González J, Lekunberri I, Álvarez-Salgado XA. (2008). Linkages between bacterioplankton community composition, heterotrophic carbon cycling and environmental conditions in a highly dynamic coastal ecosystem. *Environ. Microbiol*, 10:906-917. <https://doi.org/10.1111/j.1462-2920.2007.01509.x>
- Thompson LR, Williams GJ, Haroon MF, Shibl A, Larsen P, Shorenstein J, ... & Stingl U. (2017). Metagenomic covariation along densely sampled environmental gradients in the Red Sea. *ISME J*, 11:1, 138-151. <https://doi.org/10.3389/fmicb.2015.01438>

Thottathil SD, Balachandran KK, Gupta GVM, Madhu NV, Nair S. (2008). Influence of allochthonous input on autotrophic–heterotrophic switch-over in shallow waters of a tropical estuary (Cochin Estuary), India. *Estuar. Coast Shelf Sci.* 78(3), 551-562. <https://doi.org/10.1016/j.ecss.2008.01.018>

Thottathil SD, Balachandran KK, Jayalakshmy KV, Gupta GVM, Nair S. (2008). Tidal switch on metabolic activity: salinity induced responses on bacterioplankton metabolic capabilities in a tropical estuary. *Estuar Coastal Shelf Sci.* 78(4), 665-673. <https://doi.org/10.1016/j.ecss.2008.02.002>

Vajravelu M, Martin Y, Ayyappan S, Mayakrishnan M. (2018). Seasonal influence of physico-chemical parameters on phytoplankton diversity, community structure and abundance at Parangipettai coastal waters, Bay of Bengal, South East Coast of India. *Oceanologia*, 60(2), 114-127. <https://doi.org/10.1016/j.oceano.2017.08.003>

Wang Z, Zhang XX, Lu X, Liu B, Li Y, Long C, & Li A. (2014). Abundance and diversity of bacterial nitrifiers and denitrifiers and their functional genes in tannery wastewater treatment plants revealed by high-throughput sequencing. *PloS one*, 9(11), e113603. <https://doi.org/10.1371/journal.pone.0113603>.

Wei, T., & Simko, V. (2017). R package “corrplot”: Visualization of a Correlation Matrix (Version 0.84).

Wu DM, Dai QP, Liu XZ, Fan YP, & Wang JX. (2019). Comparison of bacterial community structure and potential functions in hypoxic and non-hypoxic zones of the Changjiang Estuary. *PloS one*, 14(6), e0217431. <https://doi.org/10.1371/journal.pone.0217431>

## Figure legends

**Fig. 1.** Station locations in the Cochin estuary (CE). A total of 8 stations were sampled. Stations S1 to S3 lies in the north of the inlet, S4 and S5 to S8 lies toward the upstream (south) of the estuary. S8 lies beyond the Thaneermukham barrage, which protects the paddy fields from salinity incursion during dry months.

**Fig. 2** Phylum level taxonomic distribution of major and minor bacterial communities from eight sampling locations during a) Wet period (b) Dry period I and (c) Dry period II, in the Cochin estuary (CE).

**Fig. 3** Genus level distribution of dominant bacterial communities, from eight sampling locations during a) Wet period (b) Dry period I and (c) Dry period II, in the Cochin estuary (CE).

**Fig. 4** Non- metric multidimensional scaling (NMDS) analysis showing different clusters of bacterial communities during a) Wet period (b) Dry period I and (c) Dry period II, in the Cochin estuary (CE).

**Fig. 5** Correlogram showing the correlations between the dominant phylum and class with environmental variables during a) Wet period (b) Dry period I and (c) Dry period II, in the Cochin estuary (CE). The Pearson correlation coefficients in the correlogram plot are colored based on the value and on the degree of association among the variables. Red and blue colors represent significant negative correlations and positive correlations. Darker color represents stronger correlations

**Fig. 6** Correlogram showing the correlations between of most dominant 20 genera with environmental variables during a) Wet period (b) Dry period I and (c) Dry period II, in the Cochin estuary (CE). The Pearson correlation coefficients in the correlogram plot are colored based on the value and on the degree of association among the variables. Red and blue colors represent significant negative correlations and positive correlations. Darker color represents stronger correlations

**Fig. S1** Monthly riverine discharge from the six major rivers draining into the Cochin estuary (CE) during the study period (July 2015 to March 2016).

**Fig. S2** Total plate count of bacteria during three seasons from eight stations in the Cochin estuary (CE).

**Fig. S3** Class level distribution of dominant bacterial phyla, Proteobacteria, Bacteroidetes and Actinobacteria from eight sampling locations during a) Wet period (b) Dry period I and (c) Dry period II, in the Cochin estuary (CE).

**Fig. S3** Canonical Correspondence Analysis (CCA) representing multiple correlation among all environmental and biological parameters. The sampling stations are marked in magenta filled circles and environmental parameters are represented as red arrows. Major bacterial phyla, class and genus are marked in black, green and blue colors, during a) Wet period (b) Dry period I and (c) Dry period II, in the Cochin estuary (CE).

**Fig. S4** Venn diagram representing the number of shared sequences between samples and unique sequences during (a) Wet period (b) Dry period I and (c) Dry period II from Cochin estuary (CE).

**Table 1** Table showing physicochemical parameters in different stations during three periods such as Wet period (Monsoon), Dry period I and (c) Dry period II in the Cochin estuary (CE).

**Table 2** This table shows the summary of the Illumina sequencing: the number of reads, % reads classified, and OTU identified, Chao-1, Shannon and Simpson species diversity are listed. Sequencing was completed for 8 samples in 3 sampling periods: Wet period (Monsoon), Dry period I and Dry period II.

Parameters	Wet Period							
	S1	S2	S3	S4	S5	S6	S7	S8
Nitrite (( $\mu$ M)	0.32	0.16	0.42	0.42	0.26	0.26	0.12	0.06
Nitrate ( $\mu$ M)	28.06	23.22	14.41	13.06	6.79	5.76	4.37	4.19
Ammonia ( $\mu$ M)	26.55	12.05	35.05	2.25	2.25	81.45	1.15	16.45
Phosphate ( $\mu$ M)	0.7	0.3	0.45	0.7	0.55	0.85	0.2	0.4
Silicate ( $\mu$ M)	115.2	119.6	61.5	15.9	86.1	138	106	101
Temp ( $^{\circ}$ C)	28.5	31.5	32.5	29	26	25	27	29
Salinity	0	0	4	16	3.6	0	0.1	0.1
DO (mL/L)	3.25	4	4.19	3.58	3.83	3.96	3.94	4.05
	Dry period I							
Nitrite (( $\mu$ M)	0.48	0.25	1.17	1.15	1.97	0.66	0.3	0.2
Nitrate ( $\mu$ M)	27.14	17.03	6.47	5.25	3.85	4.25	13.37	16.11
Ammonia ( $\mu$ M)	10.04	19.99	41.35	28.74	42.06	29.89	17.08	13.49
Phosphate ( $\mu$ M)	1.09	1.46	2.38	3.27	3.75	1.35	0.35	0.45
Silicate ( $\mu$ M)	98.71	126.5	119.7	76.39	89.74	78.9	60.95	94.9
Temp ( $^{\circ}$ C)	28.5	31.5	32.5	29	26	25	27	29
Salinity	0	4	10	26	16	13	7	5
DO (mL/L)	4.86	4.11	4.96	4.2	4.14	4.4	5.33	5.3
	Dry period II							
Nitrite (( $\mu$ M)	0.16	0.24	0.53	0.44	0.8	0.9	0.18	0.02
Nitrate ( $\mu$ M)	26.84	11.87	2.55	2.55	3.42	3.86	15.04	18.22
Ammonia ( $\mu$ M)	3.57	7.46	18.77	11.93	23.35	58.5	6.75	13.3
Phosphate ( $\mu$ M)	0.61	1.27	1.95	2.99	1.55	1.61	0.05	0.1
Silicate ( $\mu$ M)	114.83	91.19	43.01	43.01	33.9	40.27	81.55	106.3
Temp ( $^{\circ}$ C)	30.9	22.88	28.88	29.15	23.28	30.43	23.58	30.5
Salinity	0	5	21	28	19	16	6.5	2
DO (mL/L)	4.28	5.73	3.62	3.59	4.12	4.3	5.07	5.1

**Table 1** Table showing physicochemical parameters in different stations during three periods such as Wet period (Monsoon), Dry period I and (c) Dry period II in the Cochin estuary (CE).

Wet period	No of reads	% Reads, classified genus	Shannon Species Diversity	Cho-1 species Diversity	Simpson Species Diversity	Number of OTUs identified
S1	304436	75.65%	2.898	1782	0.9768	1421
S2	322227	64.49%	2.446	1798	0.9523	1324
S3	281047	65.69%	2.34	1527	0.9729	1215
S4	179025	66.01%	2.718	1388	0.9767	1121
S5	252310	64.20%	2.599	1746	0.9745	1225
S6	275375	65.91%	2.663	1720	0.977	1329
S7	236879	66.95%	2.336	1355	0.8978	948
S8	167417	64.57%	2.408	1088	0.9453	795
<b>Dry period I</b>						
S1	274745	70.93%	2.556	1930	0.9723	1543
S2	367764	67.31%	2.792	2016	0.9805	1651
S3	173542	64.60%	2.69	1600	0.9764	1203
S4	237272	71.13%	3.065	1819	0.9802	1462
S5	260109	71.13%	2.992	1879	0.9643	1575
S6	323762	66.62%	2.971	1815	0.9746	1483
S7	337895	66.74%	2.779	1887	0.9791	1558
S8	130689	70.79%	2.716	1541	0.9645	1188
<b>Dry period II</b>						
S1	171453	66.93%	2.825	1588	0.976	1078
S2	165909	52.63%	2.481	1285	0.9512	954
S3	127522	57.53%	2.239	1422	0.9706	1036
S4	189601	59.06%	2.417	1514	0.9779	1025
S5	210901	57.34%	2.41	1370	0.9769	1038
S6	176353	62.86%	2.192	1077	0.9595	829
S7	93251	63.19%	2.36	1073	0.9468	827
S8	179775	85.98%	2.376	915	0.9524	645

**Table 2** This table shows the summary of the Illumina sequencing: the number of reads, % reads classified, and OTU identified, Chao-1, Shannon and Simpson species diversity are listed. Sequencing was completed for 8 samples in 3 sampling periods: Wet period (Monsoon), Dry period I and Dry period II.




FIG 1




FIG 2






FIG 3






FIG 4

(a)



(b)



(c)






FIG 5

(a)



(b)



(c)




FIG 6




Fig S1




Fig S2




Fig S3




Fig S4




Fig S5